

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/299455884>

# Biological Control of *Botrytis cinerea* by *Bacillus* sp. Strain S7LiBe Under Abiotic Stress

Conference Paper · November 2015

CITATIONS

2

READS

460

6 authors, including:



**Leila Bensidhoum**  
Université de Béjaïa

12 PUBLICATIONS 100 CITATIONS

SEE PROFILE



**Rai Abdelwahab**  
Université de Bouira

11 PUBLICATIONS 43 CITATIONS

SEE PROFILE



**Nacera Tabli**  
Université de Béjaïa

9 PUBLICATIONS 98 CITATIONS

SEE PROFILE



**Elhafid Nabti**  
Université de Béjaïa

47 PUBLICATIONS 370 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



seaweeds in the world of pharmaceutical industry. [View project](#)



agriculture in Algeria [View project](#)

# Biological Control of *Botrytis cinerea* by *Bacillus* sp. Strain S7LiBe Under Abiotic Stress

Bensidhoum Leila<sup>\*1</sup>, Rai Abdelwahab<sup>1</sup>, Tabli Nassira<sup>1</sup>, Kahouadji Nabila<sup>1</sup>, Khaber Meriem<sup>1</sup> and Nabti Elhafid<sup>1</sup>

Department of Microbiology, University of Bejaïa, Targa Ouzemmour, Bejaïa, Algeria

## ABSTRACT

In this investigation, the isolate S7LiBe isolated from trefoil rhizosphere in Barbacha, Bejaïa (northern Algeria), was selected on the basis of plant growth promoting attributes. This isolate identified as *Bacillus* sp. based on 16S rDNA sequence analysis, exhibits the highest similarity of 99.1% with *Bacillus pichinoty* AF519463 and AF519460 and *Bacillus* sp. JX152775. *Bacillus* sp. S7LiBe has been subjected to the study of its antifungal activity against *Botrytis cinerea* *in vitro* and *in vivo* on the detached lettuce leaves. The effect of some abiotic factors on the performance of this bacterial strain as biocontrol agent and the production of some antifungal metabolites (siderophores, ammonia, hydrogen cyanide and chitinase) were also carried out. The results showed that *Bacillus* sp. S7LiBe strain inhibit the mycelial growth in high level (70%±1,33) and the fungal spores germination (82,54%±3,58). The strain produce siderophores, ammonia, hydrogen cyanide and chitinase. Its antagonistic activity was found to be optimal on CZA modified medium at a pH of 6.5 and a temperature of 30 ° C. The antagonist activity on the detached lettuce leaves showed that the bacterial strain inhibit 57.0% of the lesion extension.

**Keywords:** *Bacillus* sp., *Botrytis cinerea*, PGPR, Biocontrol.

## I. INTRODUCTION

Pathogens affecting plant health are a major menace to food production and ecosystem stability worldwide [1]. It has been estimated that approximately one third of the food crop is destroyed every year due to attack by insects, pathogenic fungi, bacteria, and nematodes [2]. *Botrytis cinerea* is one of the most hazardous plant pathogen infecting a large number of vegetable plant. This phytopathogene infect leaves, stems, flowers and fruits of plants, either by direct penetration or through wounds caused by cultivation practices [3; 4; 5; 6]. To control plant disease, producers had resort to the chemical products. However, many of the chemicals are hazardous and causes several negative effects on the environment and human health. For that reason, biocontrol has become an interesting alternative that are more "friendly" to the environment. Plant growth promoting rhizobacteria (PGPR) are the important group of microorganisms, which play a major role in the biocontrol of plant pathogens [7]. *Bacillus* species have been reported as plant promoting bacteria in a wide

range of plants [8; 9; 10; 11], it's one of the principals PGPR groups known for their application as biocontrol of several pathogenic fungi. *Bacillus* is the key bacterial genus that have shown greatest potential for *Botrytis* disease control [12], and that by several mechanisms which include; secretion of lytic enzymes, siderophores and antibiotics, competition for space and nutrient, defense plant stimulating and combination of mechanisms [7; 13]. In addition, *Bacillus* species produce spores that are resistant to heat and desiccation, which allows the preparation of more stable and durable formulations, this explains the greater availability of commercial products based on *Bacillus* [14; 15]. Commercial available biocontrol products include: Kodiak (*B. subtilis* strain GB03), Serenade (*B. subtilis* QST 713), YieldShield (*Bacillus pumilus* strain GB34), EcoGuard (*Bacillus licheniformis* strain SB3086) [16; 17]. and Subtilex (*B. subtilis* strain MB1600) which is active against *Botrytis* spp. infection of vines, strawberry, cucumber, powdery mildew of tomato, and brown rust of cereals [2; 17].

In the present investigation, *Bacillus* sp. S7LiBe strain isolated from trefoil rhizosphere in northern Algeria has been subjected to the study of its antifungal activity against *Botrytis cinerea* *in vitro* and *in vivo* on the detached lettuce leaves. Many abiotic factors, such as pH, temperature, moisture, inorganic and organic constituents, may influence the mechanisms of biocontrol, in this research we study the influence of some abiotic stress on the performance of *Bacillus* sp. S7LiBe against *B. cinerea*. The production of some antifungal metabolites (siderophores, ammonia, hydrogen cyanide and chitinase) was also carried out.

## II. METHODS AND MATERIAL

### A. Fungal strain and culture conditions

A strain *Botrytis cinerea* BC1 used in this study was obtained from the laboratory of Mycology, Bejaia University, Algeria. It was originally isolated, in the Laboratory of Plant Protection (INRA, STPV, Avignon, France), from tomato plants with typical symptoms of grey mold. The fungal strain was grown in Petri plates containing potato dextrose agar medium (PDA, Difco), The plates were incubated at 21°C for 3-10 days, Stock cultures of *B. cinerea* was maintained on PDA medium and stored at 4°C.

### B. Bacterial strain and inoculum

*Bacillus* sp. S7LiBe was isolated from trefoil rhizosphere in Barbacha, Bejaia (northern Algeria). This strain was selected for its ability to produce several promoting growth traits (production of siderophores, indole acetic acid, inorganic phosphate solubilization and heavy metal tolerance), and was identified by sequencing their entire 16S rRNA. Bacterial suspension was prepared by inoculating a loop full of cells in LB medium and incubating at 30°C for 24h. The bacterial concentration was calculated by serial dilutions and checked by optical density ( $OD_{600}$ ), and then adjusted to  $10^8$  c.f.u. /ml before use.

### C. Molecular identification

The isolate S7LiBe was subjected to the molecular identifications, genomic DNA extraction from pure bacterial colonies was carried out using the FastDNA®

SPIN kit in conjunction with the FastPrep FP120 instrument (Qbiogene, Heidelberg, Germany) according to the manufacturer's instructions. The genomic DNA was further PCR amplified for 16S-rDNA gene sequencing using the flanking primer pair 616F (5'AGA GTT TGA TYM TGG CTCAG 3') and 630R (5' CAK AAA GGA GGT GAT CC 3'). For phylogenetic analyses the obtained 16S-sequences were aligned with the *Sina Aligner V1.2.11* on the Silva website ([www.arb-silva.de](http://www.arb-silva.de)) and phylogenetical allocated with the software package ARB [18]. Phylogenetic tree construction was performed by using the Maximum-Likelihood [19].

### D. Effect of the antagonist on the growth of *B. cinerea* mycelium

The inhibitory effects of the strain on the growth of *B. cinerea* were tested by the agar diffusion method as described by [20]. 1cm<sup>2</sup> fungal plug was inoculated in the center of PDA plate, each isolate was sown at a distance of 2.5 cm from the fungus. Plates without antagonist served as control. Three replications were performed for each confrontation experiment. The plates were then incubated at 25±2°C for 5 days and verified every day. The percentage of growth inhibition (PGI) of the fungus was recorded and calculated using the formula:  $PGI (\%) = [(KR-R1)/KR] \times 100$  where KR corresponds to the distance from the point of inoculation to the colony margin on the control dish (mm). R1 represents the distance (mm) of fungal growth from the point of inoculation to the colony margin on the treated dishes.

### E. Effect of the antagonist on the germination of *B. cinerea* spores

20µl of spore suspension adjusted to  $10^6$  spores/ml and 20µl of 24h bacterial culture grown on LB medium adjusted to  $10^8$  c.f.u./ml, were pipetted into an Eppendorf tube containing 1 ml of sterile distilled water with 5% of glucose and then incubated at 21°C for 24 h. Control tubes were inoculated only with fungal spores. The experiment was realized in triplicate. The germination studies were performed on hemocytometer using a light microscope (40X).

### F. Effect of the antifungal volatiles on the growth of *B. cinerea* mycelium

The strains *Bacillus* sp. S7LiBe et *Pseudomonas* sp.S5LiBe were streaked on a Petri dish containing LB agar medium. a second Petri dish containing PDA medium, was inoculated with a 5 mm plug of the fungus *B. cinerea* in the center of the plate, and inverted and placed over the bacterial culture. The two plates were sealed together with parafilm and incubated at 25°C. The percentage of growth inhibition (n) of the fungus, was recorded and calculated using the formula:  $N = [(A-B)/A] \times 100$ , where A corresponds to the diameter of the mycelium in the control (cm) and B represents the diameter of the mycelium in the plates inoculated with bacteria (cm).

### G. In vivo antagonism experiments

The evaluation of the antagonistic activities of S7LiBe against *B. cinerea* is realized on the detached lettuce leaves. The test was done according to the protocol of INRA Avignon France with modification. Three lettuce leaves were deposited on box lined beforehand with absorbent paper soaked in distilled water, after this, the bacterial suspension ( $10^8$  CFU/ml.) was sprayed on the surface of the detached lettuce leaves. The leaves were allowed to air dry in a sterile cabinet for up to 30 min, followed by inoculation with a mycelium plug (5mm) at the center of each leaf. A control was done with lettuce leaves sprayed with sterile distilled water and then with *B. cinerea* plug in the center. The leaves were then stored in a growth chamber at 24°C for 72 h. The percentage of disease reduction of the gray mold on the lettuce leaves was calculated using the following formula:

Reduction rate (%) =  $[(A-B)/A] \times 100$ , Where; A is the lesion diameter recorded in untreated control, and B is the lesion diameter in the infected lettuce fruit treated with the antagonists [21]. For each treatment, nine leaves were assayed (three leaves as one replicate).

### H. Production of antifungal compounds

1) *HCN and NH<sub>3</sub> production*: *Bacillus* sp. S7LiBe was tested for the production of hydrogen cyanide by adapting the method of [22]. Bacteria were streaked on

nutrient agar medium amended with 4.4 g glycine/l, a Whatman filter paper no. 1 soaked in 2% sodium carbonate in 0.5% picric acid solution was placed in the top of the plate. Plates were sealed with parafilm and incubated at 28°C for 4 days. A change in color of the filter paper strip from yellow to light brown, brown or reddish brown was recorded as an indication of weak, moderate or strong production of HCN by each strain, respectively. The production of ammonia was tested in peptone water. Freshly grown cultures were inoculated into 5 ml peptone water, after incubation period 48hrs at 30°C, 0.25 ml of Nessler's reagent was added to each tube. Development of brown to yellow color indicate production of ammonia [23].

2) *Siderophores production*: The experiments were carried out in Chrome Azurol S agar according to the method of Schwyn and Neilands [24]. CAS agar was prepared from four solutions, which were sterilized separately before mixing. At 50°C, after autoclaving, nutrition solution and casamino acid solution were added to the buffer solution. Indicator solution was added last with sufficient stirring to mix the ingredients. The bacterial strain was streaked on the surface of the bleu agar plates and incubated at 30°C for 72h and examined for growth and production of orange halos surrounding the colonies.

3) *Chitinase Activity*: Chitinase activity was determined as described by Kopečný et al. [25]. Bacterial strain was inoculated in minimal salt medium containing 0.8% of colloidal chitin as carbon and energy source and incubated at 30°C. Clear zone around the colony indicate chitinase activity.

### I. Effects of culture medium, pH and temperature on the antagonistic activity of S7LiBe


The influence of the culture medium; pH; salinity and incubation temperature on the antagonistic activity of the bacterial strain was studied according to the method cited in the section (D). Four culture medium (Potato Dextrose Agar PDA; Sabouraud Dextrose Agar SDA; Czapek Dox Agar CZA; Malt Extract Agar MEA) were tested. The culture medium which gave a high PGI%, was prepared at different pH value (4.5, 6.5, 8.5, 10.5) to determine the optimum pH. The optimum temperature was studied at, 4°C; 25°C; 28°C; 30°C; 37°C and 44°C,

using the optimum medium and optimum pH. After incubation, the PGI% was calculated as described previously.

### III. RESULT AND DISCUSSION

#### A. Phylogenetic identification of S7LiBe

Comparative analysis of the 16S rDNA sequence of the strain S7LiBe with already available database showed that the Gram-positive isolate was member of the genus *Bacillus* and the 16S rRNA gene sequence was 99.1% similar to *Bacillus pichinoty* AF519463 and AF519460 and *Bacillus* sp. JX152775. The taxonomic positions of S7LiBe is shown in the phylogenetic tree (Figure 1).



**Figure 1 :** Phylogenetic relationship of S7LiBe, Dendrogram based on maximum likelihood tree calculation.


#### B. Effect of the antagonist on the growth mycelium and the germination spores of *B. cinera*

Six days after incubation, the inhibitory zones were observed in the dual culture dishes (Fig. 2a and b2). *Bacillus* sp. S7LiBe inhibited 70%± 1.33 mycelia growth of *B. cinerea* and it was effective in suppressing germination of *B. cinerea* spores (Fig. 2c and 2d). The

percentage spore germination was 25.65% ±6.69 after 24 h of incubation. These results indicate that *Bacillus* sp. S7LiBe grown on PDA plate released an extracellular diffusible metabolite(s) that inhibited the hyphal growth of *B. cinerea*. Several mechanisms have been proposed to explain the antagonist effect of *Bacillus* spp., including antibiotic production, hydrolytic enzymes synthesis, competition for nutrients, or a combination of these mechanisms in synergy [26, 27, 28]. Alabouvette et al. (2006)[29], suppose that the inhibition of spores in soil germination is due to the competition for nutrients, especially the carbon. Elad and Stewart (2007) [12] showed the sensibility of *B. cinerea* to several antifungal produced by *Bacillus* sp. *Bacillus brevis* and *Bacillus polymyxa* produce the gramicidin S and the polymyxin B which inhibit *B. cinerea* germination *in vitro* and *in vivo*.

#### C. Effect of the volatile compoundson the growth of *B. cinera mycelium*

Although the absence of direct contact between bacterial strain and the fungi, a strain *Bacillus* sp. S7LiBe inhibited up to 65.81% ±4,06 of *B. cinerea* mycelium. These result approve that *Bacillus* sp. S7LiBe release a volatile compounds which affected fungi growth. Guetsky et al. (2002)[30] showed that the volatile compounds produced by *Bacillus pumilus*, inhibit the growth of *Botrytis cinerea*. Several studies have been demonstrated that the production of volatile compounds, e.g. HCN; NH<sub>3</sub>; 2,3-butanediol; acetoin, can influence growth of fungi[31, 32] and improve the plant growth [33, 34, 35, 36].



**Figure 2 :** Antifungal activity of *Bacillus* sp. S7LiBe against *B. cinerea*.(a, c and e) *B. cinerea*BC1 Control in



agar diffusion test, spores germination test and antifungal volatiles test respectively. (b, d and f) Strain *Bacillus* sp. S7LiBe in agar diffusion test, spores germination test and antifungal volatiles test respectively.

#### D. *In vivo* antagonism experiments

*In vivo* evaluation of *Bacillus* sp. S7LiBe as antagonists towards *B. cinerea* on the lettuce leaves showed a remarkable potential (Fig. 3). The antagonist inhibited up to  $57\% \pm 1.65$  of *B. cinerea*. These results confirm those obtained in the *in vitro* tests and those reported by Wang et al. (2009)[37], they showed the potential of *Bacillus subtilis* to inhibit up to 52.4% of *B. cinerea*. According to Elad and Stewart (2007)[12], *Bo. cinerea*, is sensible to the antibiotics produced by some *Bacillus* species in several host.



**Figure 3** : Lesion extension in the lettuce leaves by *B. cinerea*. (S7) *Bacillus* sp. S7LiBe, (T) Control.

#### E. Production of antifungal compounds

1) *HCN and NH<sub>3</sub> production*: In the present study, *Bacillus* sp. S7LiBe was a moderate producer (++) of cyanide and ammonia. Researchers have reported that many *Bacillus* species could secrete several antifungal metabolites such as CHN [33, 34, 38], and NH<sub>3</sub> [34, 39]. According to Prashar et al. (2013) [38], the production of antifungal compounds diffusing in the culture medium or volatiles as cyanide and ammonia, affect negatively the fungi growth.

2) *Siderophores production*: Siderophores production is a major mechanism involved in bio-control

by many PGPR groups, including *Bacillus* sp., which produces a large variety [40]. Siderophore production in the CAS agar is characterized by the production of orange halos surrounding the colonies (Fig. 4). In this investigation, *Bacillus* sp. S7LiBe produce an important zone around the colonies and the diameter was up to  $4\text{ cm} \pm 0.6$ . Several studies have demonstrated the production of siderophores by *Bacillus* sp.[41], the role of these molecules in the control of diseases has been well documented[42], they deprive pathogenic fungi of iron since the fungal siderophores have lower affinity [43, 44]. These molecules may indirectly stimulate the biosynthesis of other antimicrobial compounds by increasing the availability of these minerals to the bacteria [45].

3) *Chitinase Activity*: It has been reported that antifungal mechanism of antagonists has been attributed to the production of hydrolytic enzymes such as chitinase, [46, 47,]. In this investigation, *Bacillus* sp. S7LiBe produce a clear zone around the colonies, indicate the production of Chitinase enzyme. Chitinase-producing microorganisms have been reported as biocontrol agents for different kinds of fungal diseases of plants [48, 49, 50]. Some studies revealed a relationship between chitinases of *Bacillus* Sp, and its ability to inhibit *Fusarium oxysporum* and *Fusarium solani* mycelial growth [51].

#### F. Effects of culture medium, pH and temperature on the antagonistic activity of S7LiBe

Understanding which environmental factors are important and how these influence disease suppression is widely recognized as a key to improving the level and reliability of biocontrol [45]. However, little knowledge of how specific factors affect the interactions among soil borne plant pathogens and their antagonists[45, 52, 53]. In this investigation, the PGI% obtained showed that *Bacillus* sp. S7LiBe inhibit the mycelial growth at different conditions tested (Fig. 4, 5 and 6), however, PDA, pH 4,5 and the temperature 4°C were the least suitable condition for *Bacillus* sp. S7LiBe ( $0,05 \geq p \geq 0,01$ ). Many factors have been discussed that may affect rhizosphere microbial communities and it is likely soil type, pH, temperature, salinity, organic carbon and inorganic nutrients, as well as the presence of other soil organisms will effect PGPR performance [54, 55, 56].

Carbon sources and minerals have been shown to have an important role in antifungal metabolite production by *Pseudomonas* BCAs, suggesting that nutrient amendments to formulations may also be a useful strategy for improving biocontrol efficacy [45].




Figure 4: Effects of culture medium on the antagonistic activity of S7LiBe




Figure 5: Effects of pH on the antagonistic activity of S7LiBe




Figure 6: Effects of temperature on the antagonistic activity of S7LiBe

#### IV. CONCLUSION

Thus the strain S7LiBe selected previously for its ability to produce several promoting growth traits, could be applied in the same time as a PGPR and a biocontrol agent, to improve crop productivity and to prevent fungal attack in crop plant.

#### V. ACKNOWLEDGMENT

Our special acknowledgments are addressed to Prof. Dr. Hartmann A. and Schmid M. from Helmholtz Zentrum München, Research Unit Microbe-Plant Interactions, Neuherberg, Germany for their molecular identification. We also want to consider this article as dedications to Prof. Dr. Hartmann for all his continuous and effective helps. We would be very grateful.

#### VI. REFERENCES

- [1] S. Compant, B. Duffy, J. Nowak, C. Clement, E. A. Barka, 2005, Use of plant growth promoting bacteria for biocontrol of plant diseases: principles, mechanisms of action, and future prospects. *Appl. Environ. Microbiol.*, 71 4951-4959.
- [2] R. Borris, Use of Plant-Associated Bacillus Strains as Biofertilizers and Biocontrol Agents in Agriculture, in *Bacteria in agrobiology: Plant growth responses*, Maheshwari DK. (Ed.), Springer, Berlin, Heidelberg 2011.
- [3] W. E. McKeen, 1974, Mode of penetration of epidermal cell walls of *vicia faba* by *botrytis cinerea*. *Phytopathology*, 64(4) 461.
- [4] M. K. Hausbeck and S. P. Penny-Packer, 1991, Influence of grower activity on concentration of airborne conidia of *Botrytis cinerea* among geranium cuttings, *Plant disease*, 75 1236-1243.
- [5] G. Peng, J. C. Sutton, D. W. Li, 1996, Sites of infection in tomato stems by *Botrytis cinerea* biological control of *Botrytis cinerea* on tomato plant by use of epiphytic yeasts *Candida guilliermondii* strain 101 and US 7 and *Candida oleophila* strain I-182: I. in vivo study *Can J. Plant Pathol Biol. Control*, 18 143-150.
- [6] T. M. O'Neil, D. Shtienberg, Y. Elad, 1997, Effect of some Host and microclimate factors on infection of tomato stems by *botrytis cinerea*, *Plant disease*, 81(1) 36-40.
- [7] Akhtar MS et Siddiqui ZA. (2010). Role of Plant Growth Promoting Rhizobacteria in Biocontrol of Plant Diseases and Sustainable Agriculture, in *Plant Growth and Health Promoting Bacteria*, Maheshwari DK. (Ed.), Springer-Verlag Berlin Heidelberg 2010.

- [8] N. Kokalis-Burelle, E. N. Vavrina, E. N. Roskopf, R. A. Shelby. 2002. Field evaluation of plant growth-promoting rhizobacteria amended transplant mixes and soil solarization for tomato and pepper production in Florida. *Plant Soil*, 238 257–266.
- [9] Y. Bai, X. Zhou, D. L. Smith. 2003. Enhanced soybean plant growth resulting from coinoculation of *Bacillus* strains with *Bradyrhizobium japonicum*. *Crop Sci* 43:1774–1781
- [10] J. W. Kloepper, C. M. Ryu, S. Zhang, 2004, Induced systemic resistance and promotion of plant growth by *Bacillus* spp. *Phytopathology* 94 1259–1266.
- [11] C. K. Deepa, S. G. Dastager, A. Pandey. 2010. Isolation and characterization of plant growth promoting bacteria from non-rhizospheric soil and their effect on cowpea (*Vigna unguiculata* (L.) Walp.) seedling growth. *World J Microbiol Biotechnol.* 26(7) 1233–1240.
- [12] Y. Elad and A. Stewart. Microbial control of *Botrytis* spp., in Y. Elad, B. Williamsom and P. Tudzynski (Eds.), *Botrytis: Biology, Pathology and Control*. Springer, Netherlands, 2007.
- [13] J. Raaijmakers, M. Vlami and T. Jorge. 2002. Antibiotic production by bacterial biocontrol agents. *Antonie van Leeuwenhoek*. 81 537-547.
- [14] M. do Vale Barreto Figueiredo, L. Seldin, F. F. de Araujo and R. de Lima Ramos Mariano. Plant Growth Promoting Rhizobacteria: Fundamentals and Applications, in Plant Growth and Health Promoting Bacteria, Maheshwari DK. (Ed.), Springer-Verlag Berlin Heidelberg 2010
- [15] O. Susana Correa and M. Abel Soria. Potential of Bacilli for Biocontrol and Its Exploitation in Sustainable Agriculture, in Plant Growth and Health Promoting Bacteria, Maheshwari DK. (Ed.), Springer-Verlag Berlin Heidelberg 2010
- [16] B. Bouizgarne. Bacteria for Plant Growth Promotion and Disease Management, in Bacteria in Agrobiolgy: Disease Management, Maheshwari DK. (Ed.), Springer-Verlag Berlin Heidelberg 2013.
- [17] V. Govindasamy, M. Senthikumar, V. Magheshwaran, U. Kumar, P. Bose, V. Sharma and K. Annapurna. *Bacillus* and *Paenibacillus* spp.: Potential PGPR for Sustainable Agriculture, in Plant Growth and Health Promoting Bacteria, Maheshwari DK. (Ed.), Springer-Verlag Berlin Heidelberg 2010.
- [18] W. Ludwig, O. Strunk, R. Westram, L. Richter, H. Meier, Yadhukumar, A. Buchner, T. Lai, S. Steppi, G. Jobb, W. Förster, I. Brettske, S. Gerber, A.W. Ginhart, O. Gross, S. Grumann, S. Hermann, R. Jost, A. König, T. Liss, R. Lüßmann, M. May, B. Nonhoff, B. Reichel, R. Strehlow, A. Stamatakis, N. Stuckmann, A. Vilbig, M. Lenke, T. Ludwig, A. Bode, K.H. Schleifer. 2004. ARB: a software environment for sequence data. *Nucl. Acids. Res.* 32 1363–1371.
- [19] G.J. Olsen, H. Matsuda, R. Hagstrom, R. Overbeek. 1994. Fast DNAm1: a tool for construction of phylogenetic trees of DNA sequences using maximum likelihood, *Computer Appl. Biosci.* 10 41-48.
- [20] I. P. Sagahón, M. A. Anducho-Reyes, H. V. Silva-Rojas, A. Arana-Cuenca, A. Alejandro Tellez-Jurado, I. O. Cárdenas-Álvarez and Y. Mercado-Flores. 2011. Isolation of Bacteria with Antifungal Activity against the Phytopathogenic Fungi *Stenocarpella maydis* and *Stenocarpella macrospora*. *International Journal of Molecular Sciences.* 12 5522-5537.
- [21] Z. N. Sadfi, I. Hannachi, D. Andurand, B. Essghaier, A. Boudabous and P. Nicot. 2008. Biological control of *Botrytis cinerea* on stem wounds with moderately halophilic bacteria. *World J Microbiol Biotechnol.* 24 2871–2877.
- [22] H. Lorck. 1948. Production of hydrocyanic acid by bacteria. *Physiol. Plant.* 1 142-146.
- [23] J. C. Cappucino and N. Sherman. 1992. Negative staining. *Microbiology: A laboratory Manual.* 3 125-179.
- [24] B. Schwyn, J. B. Neilands. 1986. Universal chemical assay for the detection and determination of siderophores, *Anal. Biochem.* 160 47-56.
- [25] J. Kopečný, B. Hodrová and C. S. Stewart. 1996. The isolation and characterization of a rumen chitinolytic bacterium. *Lett. Appl. Microbiol.* 23 195-198.
- [26] P. Marschner and S. Timonen. 2006. Bacterial community composition and activity in rhizosphere of roots colonized by arbuscular mycorrhizal fungi, in K. G. Mukerji, C. Manoharachary, J. Singh (Eds.), *Microbiam activity in the rhizosphere*, *Soil Biology.*7 139-154.
- [27] F. Jamali, A. Sharifi-Tehrani, M.P. Lutz and M. Maurhofer. 2009. Influence of Host Plant Génotype, Presence of a pathogen, and Coinoculation with *Pseudomonas fluorescens* Strains on the rhizosphere expression of hydrogen Cyanide- and 2,4-Diacetylphloroglucinol biosynthesis Genes in *P. fluorescens* biocontrol Strain CHAO. *Microb Ecol.* 57 267-275.
- [28] P. Calvo, E. Ormeño-Orrillo, E. Martínez-Romero and D. Zúñiga. 2010. Characterization of *Bacillus* isolates of potato rhizosphere from andean soils of peru and their potential PGPR characteristics. *Brazilian Journal of Microbiology.* 41 899-906.
- [29] C. Alabouvette, C. Olivain and C. Steinberg. 2006. Biological control of plant diseases. The European situation. *European Journal of Plant Pathology.* 114 329–341.
- [30] R. Guetsky, Y. Elad, D. Shtienberg et A. Dinoor. 2002. Establishment, survival and activity of the biocontrol agents *Pichia guillermondii* and *Bacillus mycoides* applied as a mixture on strawberry plants. *Biocontrol Science and Technology.* 12 705-714.
- [31] S. Alstrom. 2001. Characteristics of bacteria from oil seed rape in relation to their biocontrol of activity against *Verticillium dahliae*. *J Phytopathol.* 149 57–64.
- [32] R. E. Wheatley. 2002. The consequences of volatile organic compound mediated bacterial and fungal interactions. *Antonie Van Leeuwenhoek.* 81 357–364.



- [33] M. H. Charest, C. J. Beauchamp and H. Antoun. 2005. Effects of the humic substances of de-inking paper sludge on the antagonism between two compost bacteria and *Pythium ultimum*. *FEMS Microbiology Ecology*. 52 219–227.
- [34] O.H. Ahmed, H. Aminuddin and M. H. A. Husni. 2008. Ammonia volatilization and ammonium accumulation from urea mixed with zeolite and triple superphosphate. *Acta Agric. Scand.* 58 182-186.
- [35] P. A. Wani, M. S. Khan and A. Zaidi. 2007. Synergistic effects of the inoculation with nitrogen fixing and phosphate-solubilizing rhizobacteria on the performance of field grown chickpea. *J. Plant Nutr. Soil Sci.* 170 283-287.
- [36] C. M. Ryu, M. A. Farag, C. H. Hu, M. S. Reddy, H. X. Wei, P. W. Paré and J. W. Kloepper. 2003. Bacterial volatiles promote growth in *Arabidopsis*. *Proc. Natl. Acad. Sci.* 100 4927-4932.
- [37] S. Wang, T. Hu, Y. Jiao, J. Wie, K. Cao. 2009. Isolation and characterization of *Bacillus subtilis* EB-28, an endophytic bacterium strain displaying biocontrol activity against *Botrytis cinerea* Pers. *Front. Agric. China*. 3 247-252.
- [38] P. Prashar, N. Kapoor, and S. Sachdeva. 2013. Isolation and Characterization of *Bacillus* sp with In-vitro Antagonistic Activity against *Fusarium oxysporum* from Rhizosphere of Tomato. 15 1501-1512.
- [39] T. Kavitha, R. Nelson and J. S. Josephin. 2013. Screening of rhizobacteria for plant growth promoting traits and antifungal activity against charcoal rot pathogen *Macrophomina phaseolina*. *Int J Pharm Bio Sci.* 4 177 – 186.
- [40] E. H. Nabti, N. Mokrane, M. Ghouli, H. Manyani, M. Dary and M. G. Megias. 2013. Isolation and characterization of two halophilic *Bacillus* (*B. licheniformis* and *Bacillus* sp) with antifungal activity. *Journal of Ecology of Health & Environment*. 1 13-17.
- [41] S. H. Ji, M. A. Gururani and C. S. Chul. 2014. Isolation and characterization of plant growth promoting entophytic diazotrophic bacteria from Korean rice cultivars. *Microbiological Research*. 169 83– 98.
- [42] R. Baker, Y. Flad, B. Sneh. 1986. Physical, biological and host factors in iron competition in soils in T. R. Swinburne (ed) Iron siderophores and plant diseases. *Plenum press*, New York.
- [43] J. E. Loper and M. D. Henkels. 1999. Utilization of heterologous siderophores enhances levels of iron available to *Pseudomonas putida* in the rhizosphere. *Appl. Environ. Microbiol.* 65 5357–5363.
- [44] D. J. O’Sullivan and F. O’Gara. 1992. Traits of fluorescent *Pseudomonas* spp. involved in suppression of plant root pathogens. *Microbiol. Rev.* 56 662–676.
- [45] B. K. Duffy, G. Défago. 1999. Environmental factors modulating antibiotic and siderophore biosynthesis by *Pseudomonas fluorescens* biocontrol strains. *Applied and Environmental Microbiology*. 65 2429–38.
- [46] W. De Boer, P. J. A.K. Gunnewiek, P. Lafeber, J. D. Janse, B. E. Spit and J. W. Woldendorp. 1998. Antifungal properties of chitinolytic dune soil bacteria. *Soil Biol Biochem.* 30 193-203.
- [47] S. L. Wang, W. J. Hsiao and W. T. Chang. 2002. Purification and characterization of an antimicrobial chitinase extracellular produced by *Monascus purpureus* CCRC31499 in a shrimp and crab shell powder medium. *J Agr Food Chem.* 50 2249-55.
- [48] L. Chernin, Z. Ismailov, S. Haran, and I. Chet. 1995. Chitinolytic *Enterobacter agglomerans* antagonistic to fungal plant pathogens. *Appl. Environ. Microbiol.* 61 1720-1726.
- [49] D. Y. Kobayashi, R. M. Reedy J. A. Bick and P. V. Oudemans. 2002. Characterization of a chitinase gene from *Stenotrophomonas maltophilia* strain 34S1 and its involvement in biological control. *Appl. Environ. Microbiol.* 68 1047-1054.
- [50] S. Freeman, O. Minzm, I. Kolesnik, O. Barbul, A. Zveibil, M. Maymon, Y. Nitzani, B. Kirshner, D. Rav-David, A. Bilu, A. Dag, S. Shafir and Y. Elad. 2004. *Trichoderma* biocontrol of *Colletotrichum acutatum* and *Botrytis cinerea* and survival in strawberry. *Eur. J. Plant Pathol.* 110 361-370.
- [51] I. O. M. El-Hamshary, S. E. M. Abo-Aba, N. Awad and A. M. Gomaa. 2008. Genetic Profile of *Bacillus cereus* and *Bacillus subtilis* indigenous isolates and their performance as bio-control agent against the plant pathogen *Fusarium oxysporum*. *Research Journal of Cell and Molecular Biology*. 2 30-38.
- [52] L. L. Burpee. 1990. The influence of abiotic factors on biological control of soilborne plant pathogenic fungi. *Canadian Journal of Plant Pathology* 12 308–17.
- [53] B. H. Ownley, B. K. Duffy, D. M. Weller. 2003. Identification and manipulation of soil properties to improve the biological control performance of phenazine-producing *Pseudomonas fluorescens*. *Applied and Environmental Microbiology* 69 3333–43.
- [54] J. Van Veen, L. S. van Overbeek, J. D. van Elsas. 1997. Fate and activity of microorganisms introduced into soil. *Microbiol. Mol. Biol. Rev.* 61 121–135.
- [55] J. Sorensen. 1997. The rhizosphere as a habitat for soil microorganisms, in Modern soil microbiology, J. D. van Elsas, J. T. Trevors, E. M. H. Wellington (eds). Marcel Dekker, New York.
- [56] E. A. Paul and F. E. Clark. 1989. Soil microbiology and biochemistry. Academic Press, San Diego. p234.