



Trade Science Inc.

Environmental Science

An Indian Journal

Current Research Paper

ESAIJ, 8(3), 2013 [94-102]

Monometal and competitive sorption of heavy metals in mine soils: Influence of mine soil characteristics

Lotfi Mouni^{1*}, Lazhar Belkhiri², Djoudi Merabet³, Abdelkrim Bouzaza⁴

¹Faculté des Sciences de la Nature et de la Vie et des Sciences de la Terre, laboratoire du Gestion des ressources naturelles et assurance qualité, Université AMO de Bouira, (ALGERIA)

²Département d'hydraulique, Université de Batna 05000, (ALGERIA)

³Département de Génie des Procédés, Laboratoire de technologie des matériaux et de génie des Procédés, Université de Béjaia 06000, (ALGERIA)

⁴Equipe "Chimie et Ingénierie des Procédés" UMR6226 "Sciences Chimiques de Rennes" Ecole Nationale Supérieure de Chimie de Rennes Avenue du Général Leclerc, CS 50837, 35708 Rennes Cedex 7, (FRANCE)

E-mail : lotfimouni@gmail.com

ABSTRACT

The environmental and health effects of the contamination of soils by heavy metals depend on the ability of the soils to immobilize these contaminants. This study was conducted to assess the monometallic and competitive sorption of lead, copper, zinc and cadmium in surface samples of four soils from eastern Algeria. Sorption isotherms were characterized using Freundlich and Langmuir equations. Retention selectivity sequences indicate that, in most of the soils, Pb is the preferred retained metal, followed by Cu. The last metals in these sequences are Cd and Zn. All soils showed greater sorption capacity. On the basis of distribution coefficient values for the metal concentration of 100 mg. l⁻¹ (K_{d100}) for each soil and trace element, the two most common adsorption sequences found were Pb>>Cu>Cd>Zn and Pb>>Cu>Zn>Cd. Competition significantly reduced metal K_d , especially that of Cd. © 2013 Trade Science Inc. - INDIA

KEYWORDS

Competitive adsorption;
Soil;
Heavy metals;
Distribution coefficient.

INTRODUCTION

Assessment of the risk derived from heavy metal contamination in soils requires information about metal interaction and subsequent mobility of metals, which is directly related to their partitioning between soil and soil solution^[13]. Soils might be contaminated by heavy metals from various anthropogenic activities, including agricultural practices, transport, industrial activities, and waste disposal or mining activity. These heavy metals

are believed to be easily accumulated in the surface soil^[5,30]. Some of the elements, such as cadmium (Cd), lead (Pb), zinc (Zn), and copper (Cu) are potentially toxic towards plants and animals and can be accumulated in the food chain^[7,29]. Various soils show a very different behavior in sorption of heavy metals^[1]. This is due to the fact that the sorption capacity of each heavy metal dependent on factors such as soil pH, iron and aluminium oxide content, clay content, organic matter and cation exchange capacity^[19].

Heavy metals in the soil are to a large extent sorbed on soil particles. Adsorbed heavy metals can be desorbed by soil water, and thus move in dissolved form into plants or move to lower soil horizons and groundwater. The mobility of heavy metals in soil can be characterized by a distribution coefficient, defined as the ratio of metal concentration in the solid phase to that in liquid phase at equilibrium^[2,9-11,14,16,22]. Sorption processes and the leaching potential of trace elements by soil depend on factors such as soil pH, the nature of the mineral and organic constituents, the nature of the metal, the ionic strength of the soil solution and the simultaneous presence of competing metals^[18,23]. The adsorption of trace elements has been studied and reported in the literature for several different soil components^[3,9,10,17,21,26,31,32]. Most trace element sorption has been derived from studies conducted using single metal solutions^[28,35]. Sorption of trace elements is a competitive process between ions in solution and those sorbed onto the soil surface. In this situation, single metal solutions have limited practical applications^[14].

Batch tests were undertaken with mono and multi metal solutions with equimolar concentrations of the mentioned heavy metals in order to study the sorption equilibrium. The Langmuir isotherm model was adjusted to experimental data, allowing the determination of the soil maximum sorption capacity for each metal, with and without competition.

The present paper aims at studying the effect of competitive and monometal sorption of Pb, Cu, Cd and Zn on four soils of different physicochemical properties. The specific goals were: (1) to determine distribution coefficients K_d of these metals for a range of soils samples and investigate any correlation between distribution coefficients and soil characteristics; (2) to classify the metals and soils according to the K_d values obtained; and (3) to obtain and compare monometal and competitive solution isotherm at 20°C.

MATERIALS AND METHODS

Soil samples used in the study

Four soil samples were taken from the 20 cm deep surface horizon of Amizour, located in Bejaia, east of Algeria. The samples were transported to the laboratory in polyethylene bags, air dried and passed through

a 02 mm mesh sieve. Then they were cleaned by deionized water and dried at 50°C during 24h.

Soil analyses

Soil pH in water was determined with a pH meter using 2:1 (v:w) (water:soil) suspensions^[39]. Percentages of sand (2–0.05 mm), silt (0.05–0.002 mm) and clay (<0.002 mm) were determined by the wet sieving method and the pipette method. Exchangeable cations were extracted with 0.2M NH_4Cl ^[38] and measured by atomic absorption spectrophotometry (Ca, Mg, Na and K). Total carbon and organic carbon (C_{org}) contents in soils were determined by elemental analysis^[20]. The inorganic carbon content was calculated as the difference between the total carbon and organic carbon contents and expressed as % CaCO_3 . The X-ray diffraction (XRD) analysis of the sample was carried out via a Powder X-ray Diffractometer (BRUKERAXS) in the 2θ angle ranging from 2° to 65° employing a 0.025° step. The specific surface area was measured by N_2 gas adsorption at -196°C (COULTER-SA3100). BET equation was used to calculate the specific surface area. The physical and chemical characteristics of the samples are shown in TABLE 1.

Sorption stage

Sorption of Cu, Zn, Pb and Cd

Batch sorption studies were conducted to determine the relationship between adsorbent and adsorbate by varying the concentration of the metals. In each experiment, cadmium, copper, lead and zinc were sorbed from a “sorption solution”. The initial concentrations of Cd, Cu, Pb and Zn tested were 10, 20, 40, 50, 100 and 200 $\text{mg} \cdot \text{l}^{-1}$.

Experimental procedure: 1 g of soil sample was suspended in 100ml of sorption solution and shaken for 24 h at 20 °C in a rotary shaker. Then the suspension was centrifuged at 6000 rmin^{-1} for 5 min, and the supernatant was filtered through slow filter paper (Whatman No. 42). The supernatants metal concentrations were determined by ICP (ICP JY2000). The quantity of the metal sorbed by the soil sample was calculated by using mass balance relationship:

$$q_e = \frac{(C_0 - C_e) \times V}{W} \quad (1)$$

where q_e is the amount of adsorbed species at equilib-

Current Research Paper

rium ($\mu\text{mol. g}^{-1}$), C_0 is the initial concentration of the species in solution ($\mu\text{mol. l}^{-1}$), C_e is the measured concentration ($\mu\text{mol. l}^{-1}$) of metal in solution when the equilibrium is reached. V is the solution volume (L) and W is the weight of air-dried soil (g).

The pH was not controlled during the experiments; it was monitored before filtration and analysis (after equilibrium).

Adsorption from mixtures

The experimental procedure was the same as the monocomponent adsorption (see above). 1 g of soil was mixed with 100 ml of a solution containing copper, zinc, lead and cadmium in 1:1 mg ratio (total concentration range 10–100 mg l^{-1}).

Adsorption isotherms

The equilibrium data were analyzed in accordance with the Langmuir and Freundlich sorption isotherms. These isotherms allow describing adsorption phenomena of metals from aqueous solution onto soil sample. The Langmuir isotherm is expressed by:

$$C_e/q_e = 1/(bqm) + C_e/qm \quad (2)$$

where q_e and C_e are the adsorption capacity and concentration, respectively, at the equilibrium. qm is the maximum adsorption capacity at saturation ($\mu\text{mol metal g}^{-1}$) and b is the affinity coefficient ($\mu\text{mol. l}^{-1}$) (Langmuir, 1918), q_e can be expressed as:

$$q_e = K_d.C_e \quad (3)$$

where K_d is the distribution coefficient that characterizes the affinity of the metal for the sorbent, the Freundlich isotherm, which assumes that different sites with several adsorption energies are involved, is expressed by:

$$q_e = K_f.C_e^{1/n} \quad (4)$$

where K_f (lk g^{-1}) is the Freundlich distribution coefficient related to the total adsorption capacity of the solid, and n is the Freundlich sorption exponent.

The goodness-of-fit for all equations was estimated by the regression coefficient. Statistical analysis was performed using the Excel program. In addition, for each stage of each experiment, the distribution of each metal i between soil and solution following equilibration was expressed in terms of the quotient:

$$K_{d,i} = \frac{C_{i,\text{soil}}}{C_{i,\text{solu}}}$$

where $C_{i,\text{soil}}$ is the concentration of metal i on the soil (mmol kg^{-1}) and $C_{i,\text{solu}}$ is the concentration of metal i in solution (mmol l^{-1})^[3,31]. The distribution coefficient at 100mg l^{-1} ($K_{d_{100}}$) according to Covelo et al. 2007 was also used to evaluate capacity of soils to adsorb heavy metals.

RESULTS AND DISCUSSION

Soil characteristics

TABLE 1 summarizes the main soil sample characteristics. The four soils were basic (pH 7.5–8.3). pH is one of the main factors controlling metal adsorption, mobility and availability of metals. The organic carbon (Corg) content ranging from 0.96 to 1.34%. The Soil 2 had a highest cation exchange capacity (CEC). Soil 2, Soil 3 and Soil 4 have a clay texture, but Soil 1 contains a higher proportion of silt. On the other hand, the soils have a similar surface area except for Soil 1 which has the lowest surface area.

TABLE 1 : Main characteristics of soil samples

| Soil property | soil1 | soil2 | soil3 | soil4 |
|---|-------|-------|-------|-------|
| Clay (%) | 24.6 | 60.2 | 51.9 | 54.7 |
| Sand (%) | 28.1 | 11.1 | 7.9 | 10.1 |
| Silt (%) | 47.3 | 28.7 | 40.2 | 35.2 |
| pH | 7.5 | 7.9 | 8.3 | 7.8 |
| Corg | 1.34 | 1.12 | 1.08 | 0.96 |
| CEC cmoles kg^{-1} | 13.8 | 46.3 | 34.6 | 37.6 |
| Surface Area ($\text{m}^2 \text{g}^{-1}$) | 12.1 | 41.8 | 22.3 | 29.9 |
| Goethite (%) | 28.7 | 32.2 | 47.3 | 18.32 |
| Quartz (%) | 3.4 | 2.65 | 1.74 | 1.24 |
| Feldspars (%) | 7.2 | nd | n. d | nd |
| Kaolinite (%) | 53.4 | 24.5 | 31.4 | 23.8 |
| Mica (%) | 7.7 | 5.5 | 12.2 | 12.1 |
| Smectite clay content (%) | 28.1 | 67.3 | 54.6 | 62.1 |

Adsorption isotherms and K_d values

The adsorption isotherms of single metal ions onto soil samples are reported in Figure 1. These isotherms can be subdivided in three parts. At low-concentration, the metal introduced is adsorbed proportionally. At intermediate, there is a pseudo-plateau, which probably corresponds to the saturation of the surface. However in the case of lead, we note that the amount adsorbed increases regularly when the solution concentration




Figure 1 : Monometal sorption isotherms at 20°C for retention of Pb, Cu, Cd and Zn by four soils (soil1, soil2, soil3 and soil4).

increases. This is probably due to a surface precipitation of lead.

TABLE 2 shows values of parameters deduced from the application of Langmuir and Freundlich equa-




Figure 2 : Competitive sorption isotherms at 20°C for retention of Pb, Cu, Cd and Zn by four soils (soil1, soil2, soil3 and soil4).

tions. The maximum adsorption capacity (q_m), deduced from the Langmuir equation, may be useful in compar-

Current Research Paper

ing the potential adsorption capacity for the studied soils. Langmuir sorption parameters of the studied soils showed differences among the studied heavy metals, as reflected by their sorption maxima (q_m) and bonding energy coefficient (b) (TABLE 2). Among all studied metals, Cu showed the highest value of adsorption maxima (q_m) in all studied soils. Much lower value of adsorption maxima were obtained for Cd. In addition, the bonding energy coefficients (b) for Cd and Zn were lower than those for Cu, suggesting that Cd and Zn could not form much stronger complexes than Cu. Moreover, we note that the data of Pb follow C isotherm. In the other hand, the values for the Freundlich distribution (K_f) coefficient were 9.46 and 2.53 litre kg^{-1} for Zn and Pb sorption by Soil 2, respectively. Concerning n parameter, which expresses site heterogeneity, it varied within a wide range of values (0.27–4.83) for the various metal and soil. The amount of Pb and Cu sorbed was greater than that of Zn and Cd.

TABLE 2 : Isotherm parameters of four soils deduced from monometal sorption in solution at 20°C.

| Soil | Isotherms | | | | | |
|-------|------------------------|----------------------|-------|----------------|-------|-------|
| | Langmuir | | | Freundlich | | |
| | $Q_m (\mu mol g^{-1})$ | $b (L \mu mol^{-1})$ | R^2 | $K_f (l kg l)$ | n | R^2 |
| Zn | | | | | | |
| Soil1 | 50 | 0.0062 | 0.99 | 3.32 | 2.74 | 0.87 |
| Soil2 | 125 | 0.0072 | 0.98 | 9.46 | 2.99 | 0.99 |
| Soil3 | 111.11 | 0.0056 | 0.96 | 9.15 | 3.24 | 0.98 |
| Soil4 | 166.66 | 0.0089 | 0.97 | 12.44 | 2.98 | 0.99 |
| Cd | | | | | | |
| Soil1 | 32.26 | 0.0148 | 0.99 | 4.08 | 3.33 | 0.92 |
| Soil2 | 100 | 0.010 | 0.75 | 8.65 | 2.94 | 0.87 |
| Soil3 | 66.66 | 0.0098 | 0.80 | 7.25 | 3.3 | 0.89 |
| Soil4 | 100 | 0.012 | 0.94 | 7.65 | 2.77 | 0.98 |
| Cu | | | | | | |
| Soil1 | 90.9 | 0.02 | 0.99 | 19.45 | 4.83 | 0.75 |
| Soil2 | 166.66 | 0.04 | 0.99 | 32.26 | 4.21 | 0.88 |
| Soil3 | 166.66 | 0.038 | 0.99 | 28.27 | 4.06 | 0.89 |
| Soil4 | 200 | 0.039 | 0.97 | 36.23 | 4.098 | 0.78 |
| Pb | | | | | | |
| Soil1 | 142.85 | 0.016 | 0.99 | 4.92 | 1.68 | 0.91 |
| Soil2 | a | a | a | 2.53 | 0.67 | 0.77 |
| Soil3 | a | a | a | 6.72 | 1.03 | 0.55 |
| Soil4 | a | a | a | 0.074 | 0.27 | 0.76 |

^a : Langmuir equation did not fit to the data.

The adsorption data were fitted better by the Langmuir equation than by the Freundlich equation. Distribution coefficients (K_d) represent the affinity of the metallic cations in solution towards the solid phase. They can be used to evaluate the sorption and retention of the metallic cations in soils^[11]. A distribution coefficient can be related to both plant uptake and environmental pollution.

The low values of distribution coefficients indicate

TABLE 3 : Values of K_d for sorption in four soils deduced from monometal sorption in solution at 20°C.

| Initial Concentration (mg/l) | Soil1 | Soil2 | Soil3 | Soil4 |
|------------------------------|-------|-------|-------|-------|
| Zn | | | | |
| 08.5 | 282 | 6000 | 5237 | 10575 |
| 16.3 | 211 | 1200 | 1006 | 3019 |
| 26.9 | 115 | 668 | 391 | 1322 |
| 33.3 | 117 | 453 | 324 | 1302 |
| 44.6 | 125 | 365 | 291 | 488 |
| 90.9 | 43 | 138 | 86 | 208 |
| 185.8 | 19 | 66 | 55 | 100 |
| Cd | | | | |
| 09.1 | 647 | 5973 | 3041 | 5625 |
| 18 | 310 | 3500 | 2276 | 3000 |
| 27.2 | 210 | 2416 | 928 | 1068 |
| 36.5 | 162 | 1307 | 563 | 934 |
| 44.9 | 137 | 187 | 206 | 466 |
| 90 | 45 | 280 | 74 | 185 |
| 182.6 | 23 | 142 | 81 | 121 |
| Cu | | | | |
| 08.86 | 4190 | 49111 | 44200 | 52058 |
| 17.6 | 11600 | 10294 | 18041 | 24718 |
| 27.65 | 2557 | 21153 | 23000 | 61400 |
| 36.45 | 2447 | 15739 | 15083 | 37142 |
| 45.67 | 407 | 6818 | 3112 | 37916 |
| 90.68 | 90 | 346 | 328 | 448 |
| 184.52 | 43 | 124 | 101 | 197 |
| Pb | | | | |
| 09.84 | 1934 | 2381 | 2333 | 2333 |
| 19.61 | 1615 | 4360 | 3533 | 2987 |
| 30.94 | 2108 | 6382 | 6565 | 5477 |
| 40.34 | 1666 | 6198 | 7870 | 7631 |
| 50.39 | 1516 | 7153 | 9524 | 9138 |
| 101.89 | 1221 | 15007 | 19403 | 13567 |
| 198.01 | 320 | 8376 | 5211 | 23963 |

that most of the metals present in the system remain in the solution and are available for transport, chemical processes, and plant uptake^[27]; whereas higher values indicate lower mobility and higher retention of metals in the soil. Therefore, to be able to compare the different metal cations in each different soil, based on the reasoning of Covelo et al 2007, a K_d at a given concentration is considered.

The most used value is K_{d100} , which represents the ratio of metal sorbed to equilibrium concentration at 100 mg l⁻¹ metal in the added solution^[11]. TABLE 4 shows the value of K_{d100} for all soils. The K_d values (TABLE 3) obtained for each metal for the concentrations and soils tested (in 1 kg⁻¹) are as follows: Cd (23–5973), Cu (43–37916), Pb (320–23963) and Zn (19–6000). Zinc showed the lowest K_d values regardless of the soil sample, while Pb K_d were generally the highest. The obtained K_d sequence for both metals showed good agreement with heavy metal K_d data in soils compiled from the literature^[9,10,32]. The lowest K_d values were obtained for the Soil1, which correspond to the sample with the highest sand fraction and also the lowest CEC. This can be related to the fact that distribution of trace elements in polluted soils is known to follow particle size distribution^[17]. The lowest sorption capacity is related to the sand fraction, due to the lower surface area associated with larger particle size fractions^[21]. On the contrary, the highest K_d values corresponded to Soil2, Soil3 and Soil4 for all metals. This is due to the high clay fraction^[17]. It means that Zn and Cd have the highest mobility associated to the lowest adsorption. In contrast Cu and Pb present the opposite behaviour. The K_d quantification was extremely dependent on the initial metal concentration. Thus we note a decreases of the K_d values, especially for Cu and Zn, when the initial metal concentration was increased^[26]. This decrease reflects the decreasing average binding energy of occupied sites as concentration increases. The sites occupied at high concentration including both high- and low-energy sites, whereas those occupied at low concentration are all of relatively high energy^[15]. The higher K_d value that was obtained in the experiment with lower metal concentrations is associated with the sorption sites of high selectivity, which have relatively strong bonding energies^[6,33].

TABLE 4 : Values of K_{d100} (Lkg⁻¹) for the different soils and selectivity sequences deduced from monometal sorption in solution at 20°C.

| | Pb | Cu | Cd | Zn | Selectivity sequences |
|-------|-------|-----|-----|-----|-----------------------------|
| Soil1 | 1222 | 90 | 45 | 43 | Pb>Cu>Cd?Zn |
| Soil2 | 15007 | 346 | 280 | 138 | Pb>>Cu>Cd>Zn |
| Soil3 | 19404 | 329 | 74 | 87 | Pb>>Cu>Zn>Cd |
| Soil4 | 13567 | 448 | 186 | 209 | Pb>>Cu>Zn>Cd |
| Pb | | | | | Soil3> Soil4> Soil2> Soil1 |
| Cu | | | | | Soil4> Soil2> Soil3>>Soil1 |
| Cd | | | | | Soil2> Soil4>> Soil3> Soil1 |
| Zn | | | | | Soil4> Soil2> Soil3> Soil1 |

The most sorbent and retentive soil for the set of heavy metals was Soil4 followed by Soil 2 and Soil 3. The least sorbent and retentive soil was Soil 1. This indicates that the highest adsorption occurs for Soils that have a high cation exchange capacity values (CEC). The lowest adsorption was observed for soil 1 that contains low amount of clay and a low cation exchange capacity value.

According to the K_{d100} values, the two adsorption sequences most found were Pb>>Cu>Zn>Cd and Pb>>Cu>Cd>Zn. These sequences follow approximately the order of the electronegativity of the metal cations, which is, Pb (2.3) Cu (1.9), Cd (1.7) and Zn (1.6). The electronegativity is an important factor in determining which of the trace metals will chemisorb with the highest preference^[25]. These results are in agreement with that found by Arias et al^[4]. The adsorption of Pb and Cu, by acid soils, is higher than that of Cd, Ni and Zn. The higher affinity of the four soils for Pb is probably due to the existence of a greater number of active sites (mostly organic matter) with high specificity. When Pb is present these sites would not be occupied by other cations. This affinity can be attributed to: (i) its greater first hydrolysis constant, and therefore, it's more readily hydrolyzed, (ii) its higher atomic weight, (iii) its higher ionic radius, and subsequently smaller hydrated radius, and (iv) its larger Misono softness value, making it a better candidate than other metals for electrostatic and inner-sphere surface complexation reactions. According to the sequences presented by Schwertmann and Taylor 1989, Zn is always adsorbed to a larger extent than Cd on the synthetic samples, which was the case for our study.

Current Research Paper

TABLE 5 : Isotherm parameters of four soils deduced from competitive sorption in solution at 20°C.

| Soil | Isotherms | | | | | |
|-------|------------------------------|---------------------------|-------|--------------------------|-------|-------|
| | Langmuir | | | Frendlich | | |
| | $Q_m (\mu\text{mol g}^{-1})$ | $b (L\mu\text{mol}^{-1})$ | R^2 | $K_f (l\text{ kg}^{-1})$ | n | R^2 |
| Zn | | | | | | |
| Soil1 | 0.43 | -0.0037 | 0.84 | 343.73 | -1.37 | 0.50 |
| Soil2 | 35.71 | 0.008 | 0.96 | 07.55 | 5.26 | 0.84 |
| Soil3 | 25 | 0.0011 | 0.96 | 07.34 | 6.33 | 0.62 |
| Soil4 | 22.72 | 0.0178 | 0.99 | 08.71 | 8.00 | 0.84 |
| Cd | | | | | | |
| Soil1 | 5.31 | 0.015 | 0.81 | 9.12 | -7.75 | 0.20 |
| Soil2 | 14.29 | 0.029 | 0.98 | 06.02 | 8.26 | 0.79 |
| Soil3 | 09.25 | 0.056 | 0.98 | 05.11 | 11.11 | 0.51 |
| Soil4 | 10.63 | 0.091 | 0.99 | 06.46 | 12.82 | 0.76 |
| Cu | | | | | | |
| Soil1 | 55.55 | 0.032 | 0.99 | 09.07 | 4.56 | 0.89 |
| Soil2 | 125 | 0.015 | 0.95 | 14.16 | 3.35 | 0.92 |
| Soil3 | 83.33 | 0.026 | 0.98 | 17.02 | 4.38 | 0.88 |
| Soil4 | 100 | 0.026 | 0.99 | 13.26 | 3.21 | 0.86 |
| Pb | | | | | | |
| Soil1 | 33.33 | 0.05 | 0.99 | 3.82 | 2.32 | 0.87 |
| Soil2 | 55.55 | 0.042 | 0.91 | 3.69 | 1.65 | 0.8 |
| Soil3 | 38.45 | 0.05 | 0.99 | 3.92 | 2.14 | 0.83 |
| Soil4 | 50 | 0.045 | 0.96 | 4.075 | 1.9 | 0.76 |

^a : Langmuir and Freundlich equations did not fit to the data.

TABLE 7 : Desorption ($\mu\text{mol g}^{-1}$) of Pb and Zn into NaNO_3 solution from soils previously loaded by equilibration in solutions with various concentrations of the corresponding metal ($\mu\text{mol l}^{-1}$) values shown are means \pm standard deviations.

| Soil | Initial concentration of metal in loading solution ($\mu\text{mol l}^{-1}$) | | | | | | | | | |
|--------|---|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| | 700 | | 1200 | | 1600 | | 2300 | | 3300 | |
| | Pb | Zn | Pb | Zn | Pb | Zn | Pb | Zn | Pb | Zn |
| | $\mu\text{mol g}^{-1}$ | | | | | | | | | |
| Soil 1 | 2.2 \pm 0.03 | 6.4 \pm 0.01 | 2.3 \pm 0.05 | 7.2 \pm 0.03 | 3.5 \pm 0.03 | 7.8 \pm 0.00 | 4.5 \pm 0.01 | 8.1 \pm 0.03 | 5.8 \pm 0.06 | 9.2 \pm 0.01 |
| Soil 2 | 3.1 \pm 0.05 | 6.8 \pm 0.03 | 3.6 \pm 0.03 | 7.5 \pm 0.01 | 3.8 \pm 0.02 | 7.4 \pm 0.03 | 4.7 \pm 0.02 | 8.3 \pm 0.03 | 5.9 \pm 0.04 | 9.6 \pm 0.02 |
| Soil 3 | 3.8 \pm 0.03 | 6.3 \pm 0.05 | 3.3 \pm 0.03 | 7.1 \pm 0.06 | 3.9 \pm 0.02 | 7.6 \pm 0.05 | 5.1 \pm 0.04 | 8.6 \pm 0.06 | 5.3 \pm 0.02 | 9.0 \pm 0.03 |

very strong. Although competition reduced sorption of all four metals, the magnitude of these effects was different for each metal. Thus the effect of competition in reducing the sorption of metals followed generally the order Pb>Cu>Cd>Zn.

In soils with high pH the predominance of the metals is in the hydrolyzed (MOH^+) form^[16]. In this case, the influence of the metal hydrolysis on metal adsorption becomes more important for the more alkaline range of pH, as suggested by Elliot et al^[12], Brümmer et

TABLE 6 : Values of Kd_{100} (L kg^{-1}) for the different soils and selectivity sequences deduced from competitive sorption in solution at 20°C.

| | Pb | Cu | Cd | Zn | Selectivity sequences |
|-------|-----|-------|------|-------|-----------------------------|
| Soil1 | 157 | 54.6 | 7.5 | 0.32 | Pb>Cu>Cd>Zn |
| Soil2 | 537 | 178.3 | 21.2 | 28.2 | Pb>>Cu>Zn>Cd |
| Soil3 | 213 | 91.8 | 13.0 | 19.06 | Pb>>Cu>Zn>Cd |
| Soil4 | 368 | 133.8 | 15.3 | 17.6 | Pb>>Cu>Zn>Cd |
| Pb | | | | | Soil2> Soil4> Soil3> Soil1 |
| Cu | | | | | Soil2> Soil4> Soil3>>Soil1 |
| Cd | | | | | Soil2> Soil4>> Soil3> Soil1 |
| Zn | | | | | Soil2> Soil3> Soil4> Soil1 |

Effect of competition on metal adsorption

Comparison of monometal and competitive sorption isotherm parameters (TABLES 2 and 5) reveals that the presence of other cations decreases sorption of all metals. In fact, competition among the four metals reduced their Kd_{100} values significantly for all the soils studied (TABLE 6). This suggests that the metals were competing for the same binding sites. Although Kd_{100} values indicated a reduction in metal adsorption due to competition, metal isotherms (Figure 1) show that the differences in metal sorption were evident mostly at the higher end of equilibrium concentrations. Thus at low metal concentrations, effects of competition were not

al. (1988), and Schwertmann and Taylor (1989). In general, metal characteristics such as the order of the electronegativity^[25], the charge-to-radius ratio^[16] or metal-ion hydrolysis constants^[40], sequences do not always explain metal bonding selectivity to heterogeneous soil systems^[35]. Chemical inputs to soils can take several pathways, including rapid leaching into groundwaters, uptake by plants, volatilization to the atmosphere, and storage and retention by soil.

Desorption

Depending on the concentrations of the solutions from which initial adsorption had occurred, desorption into NaNO_3 solution ranged from 2.2 to 5.9 $\mu\text{mol g}^{-1}$ for Pb and from 6.1 to 9.8 $\mu\text{mol g}^{-1}$ for Zn, and for a given concentration of the initial adsorption solution was always greater for Zn than for Pb (TABLE 7). The slight increase in desorption with the concentration of the initial adsorption solution that was observed in this study is attributable to the increased occupancy of binding sites, which reduces binding energy.

ACKNOWLEDGEMENTS

This work was supported by the University of A.Mira of Bejaia, Algeria. The authors thank Dr. E. Guibal for essential comments and enjoyable discussion in the organization of the research and Jean Marie Toulomes for skillful laboratory assistance.

CONCLUSION

A detailed investigation of single and competitive sorption processes was performed in four soils. Most sorption isotherms for trace elements were adequately described by the Langmuir equation. The sorption capacity of the soils for Cu, as measured by the estimated q parameter from the Langmuir equation was greater than for other trace elements. K_d values in monometal followed the two adsorption sequences $\text{Pb} \gg \text{Cu} > \text{Zn} > \text{Cd}$ and $\text{Pb} \gg \text{Cu} > \text{Cd} > \text{Zn}$. These sequences follow approximately the order of the electronegativity of the metal cations. When the studied metals were in competition, this order did not change, although the K_d values of all metals decreased significantly. In general, Cu and Pb were sorbed in greater amounts with binding strength compared to Zn and Cd. This suggests that sorption does not only depend on the element, but also on the soil reactive phase and on the element concentration. The generally greater distribution coefficient (e.g., K_{d100}) led to better retention of trace elements against leaching.

REFERENCES

- [1] P.Alumaa, U.Kirso, V.Petersell, E.Steinnes; Sorption of toxic heavy metals to soil, International Journal of Hygiene and Environmental Health, **204(5-6)**, 375-376 (2002).
- [2] B.J.Alloway; Heavy Metals in Soils, Blackie Academic and Professional, London, (1995).
- [3] R.Anderson, T.H.Christensen; Distribution Coefficients of Cd, Co, Ni and Zn in Soils, J.Soil.Sci., **39**, 15-22 (1988).
- [4] M.Arias, C.Pérez-Novo, F.Osorio, E.López, B.Soto; Adsorption and desorption of copper and zinc in the surface layer, of acid soils, Journal of Colloid and Interface Science, **288**, 21-29 (2005).
- [5] D.L.Baker; Copper.In, Alloway BJ (Ed) Heavy metals in soils, Blackie & Sons, London, 151-176 (1990).
- [6] N.T.Basta, M.A.Tabatabai; Effect of cropping systems on adsorption of metals by soils, III.Competitive adsorption, Soil.Sci., **153**, 331-337 (1992a).
- [7] G.Berndes, F.Fredrikson, P.Börjesson; Cadmium accumulation and Salix-based phytoextraction on arable land in Sweden, Agriculture, Ecosystems and Environment, **103**, 207-223 (2004).
- [8] G.W.Brummer; in: M.Bernhard, F.E.Brinckman, P.J.Sadler, (Eds), The Importance of Chemical Speciation in Environmental Processes, Springer, Berlin, 169 (1986).
- [9] E.F.Covelo, M.L. Andrade, F.A.Vega, Competitive sorption and desorption of heavy metals by individual soil components, Journal of Hazardous Materials, **140**, 308-315 (2007).
- [10] E.F.Covelo, F.A.Vega, M.L.Andrade, Simultaneous sorption and desorption of Cd, Cr, Cu, Ni, Pb, and Zn in acid soils I, Selectivity sequences, Journal of Hazardous Materials, **147**, 852-861 (2007).
- [11] E.F.Covelo, M.L.Andrade, F.A.Vega, Competitive adsorption and desorption of cadmium, chromium, copper, nickel, lead and zinc by Humic Umbrisols, Comm, Soil.Sci.Plant Anal, **35(19/20)**, 2709-2729 (2004).
- [12] H.A.Elliot, M.R.Liberati, C.P.Haung, Competitive adsorption of heavy metals by soils, J.Environ.Qual., **15**, 214-217 (1986).
- [13] L.J.Evans; Chemistry of metal retention by soils, Several processes are explained, Environ.Sci. Technol., **23**, 1046-1056 (1989).
- [14] M.P.F.Fontes, A.T.de Matos, L.M.da Costa, J.C.L.Neves, Competitive adsorption of Zn, Cd, Cu

Current Research Paper

- and Pb in three highly weathered Brazilian soils, *Commun.Soil.Sci.Plant Anal.*, **31**, 2939-2958 (2000).
- [15] C.W.Gray, R.G.McLaren, A.H.C.Roberts, L.M.Condron; Cadmium phytoavailability in some New Zealand soils, *Aust.J.Soil.Res.*, **37**, 461-477 (1999).
- [16] P.C.Gomes, M.P.F.Fontes, D.G.da Silva, E.de, S.Mendonc, A.R.Netto; Selectivity sequence and competitive adsorption of heavy metals by Brazilian soils, *Soil.Sci.Soc.Am.J.*, **65**, 1115-1121 (2001).
- [17] Z.Hartyáni, E.Dávid, S.Szabó, V.Szilágyi, T.Horváth, A.H.Tóth; Determination of the trace elements distribution of polluted soils in Hungary by X-ray methods, *Microchem.J.*, **67**, 195-200 (2000).
- [18] R.D.Harter, R.Naidu; An assessment of environmental and solution parameter impact on trace-metal sorption by soils, *Soil.Sci.Soc.Am.J.*, **65**, 597-612 (2001).
- [19] L.Hernandez, A.Probst, J.Probst, E.Ulrich; Heavy metals distribution in some French forest soils, Evidence for atmospheric contamination, *Science of the Total Environment*, **312**, 195-219 (2003).
- [20] ISO, Soil quality, 1996, Determination of organic and total carbon after dry combustion (elementary analysis) ISO, Analysis, Part 3, SSSA Book Ser.5, SSSA, Madison, WI, **10694**, 475-490 (1995a).
- [21] C.K.Jain, D.Ram; Adsorption of lead and zinc on bed sediments of the river Kali, *Water.Res.*, **31**(1), 154-62 (1997).
- [22] A.D.Karathanasis; Subsurface migration of copper and zinc mediated by soil colloids, *Soil.Sci.Soc. Am.J.*, **63**, 830-838 (1999).
- [23] R.S.Kookama, R.Naidu; Effect of soil solution composition on cadmium transport through variable charge soils, *Geoderma*, **84**, 235-248 (1998).
- [24] I.Langmuir; The adsorption of gases on plane surface of glass, mica and platinum, *J.Am.Chem.Soc.*, **40**, 1361-1403 (1918).
- [25] M.B.McBride; *Environmental Chemistry of Soils*. Oxford University Press, New York, (1994).
- [26] Mesquita and Vieira e Silva; Zinc adsorption by a calcareous soil, Copper interaction *Geoderma*, **69**, 137-146 (1996).
- [27] M.T.Morera, J.C.Echeverria, C.Mazkarian, J.J.Garrido, Isotherm and sequential extraction procedures for evaluating sorption and distribution of heavy metals in soils, *Environ.Pollut.*, **113**, 135-144 (2001).
- [28] A.A.Ponizovsky, E.H.Allen, J.A.Ackerman; Copper activity in soil solutions of calcareous soils, *Environ.Pollut.*, **10**, 1-6 (2006).
- [29] F.Qin, X.Shan, B.Weil; Effects of low-molecular-weight organic acids and residence time on desorption of Cu, Cd, and Pb from soils *Chemosphere*, **57**, 253-263 (2004).
- [30] L.Samsøe-Petersen, E.H.Larsen, P.B.Larsen, P.Bruun; Uptake of trace elements and PAHs by fruit and vegetable from contaminated soils *Environ Sci.Technol.*, **36**, 3057-3063 (2002).
- [31] M.J.Sánchez-Martin, M.Sánchez-Camazano; Adsorption and mobility of cadmium in natural, uncultivated soils, *J.Environ.Qual.*, **22**, 737-742 (1993).
- [32] J.Sastre, G.Rauret, M.Vidal; Sorption-desorption tests to assess the risk derived from metal contamination in mineral and organic soils, *Environment International*, **33**, 246-256 (2007).
- [33] J.Sastre, G.Rauret, M.Vedal; Effect of the cationic composition of sorption solution on the quantification of sorption-desorption parameters of heavy metals in soils, *Environ.Pollut.*, **140**, 322-339 (2006).
- [34] U.Schwertmann, R.M.Taylor; Iron oxides, In: J.B.Dixon, S.B.Weed, (Ed); *Minerals in soil environments*, ASA and SSSA, Madison, WI, 379-438 (1989).
- [35] S.Serrano, F.Garrido, C.G.Campbell, M.T.Garcia-Gonzalez; Competitive sorption of cadmium and lead in acid soils of Central Spain, *Geoderma*, **124**, 91-104 (2005).
- [36] G.Sposito; *The Chemistry of Soils*, Oxford Univ.Press, New York, (1989).
- [37] F.J.Stevenson, A.Fitch; Chemistry of complexation of metal ions with soil solution organics, In: P.M.Huang, et al, (Ed); *Interactions of soil minerals with natural organics and microbes*, ASA and SSSA, Madison, WI, 29-58 (1986).
- [38] M.E.Sumner, W.P.Miller; Cation exchange capacity and exchange coefficients, *Methods of Soil Analysis, Part 3, Chemical Methods*, SSSA Book Series, D.L.Sparks, Madison, Wisconsin, USA, **5**, (1996).
- [39] G.W.Thomas; Soil pH and soil acidity, in: D.L.Sparks, A.L.Page, P.A.Helmke, R.H.Loeppert, P.N.Soltanpour, M.A.Tabatabai, C.T.Johnson, M.E.Sumner; *Methods of Soil analysis, Part3-chemical methods*, (1996).
- [40] G.Welp, G.W.Brummer, Adsorption of solubility of ten metals in soil samples of different composition, *J.Plant.Nutr.Soil.Sci.*, **162**, 155-161 (1999).