

# Phytochemical analysis of Myrtus communis plant: Conventional versus microwave assisted-extraction procedures

Auteurs

Nadia Bouaoudia-Madi, Lila Boulekbache-Makhlouf, Nabil Kadri, Farid Dahmoune, Hocine Remini, Sofiane Dairi, Sonia Oukhmanou-Bensidhoum, Khodir Madani

Date de publication

2017/6/10

Revue

Journal of Complementary and Integrative Medicine

Volume

14

Numéro

4

Éditeur

De Gruyter

Description:

Background

Myrtle (*Myrtus communis* L) may constitute an interesting dietary source of health protective compounds. Microwave-assisted extraction (MAE) of total phenolic compounds (TPC) from myrtle leaf, stems, pericarp, and seeds was studied and the results were compared with those of the conventional method extraction (CME) in terms of extraction time.

Methods

Extraction yield/efficiency and antioxidant activity were measured using radical scavenging assay (DPPH•) and reducing power.

Results

The results show that the MAE was higher in terms of saving energy, extraction time (62 s) and extraction efficiency of bioactive compound compared to CME (2 h). Leaf presented the optimum content of total phenols (250 mg GAE. g<sup>-1</sup> DW) and flavonoids (13.65 mg GAE. g<sup>-1</sup> DW). However, the anthocyanin content was most important in pericarp extract (176.50±2.17 mg Cyd-3-glu g<sup>-1</sup> DW). The antioxidant activity ...

Nombre total de citations

Cité 5 fois

Articles Google Scholar

Phytochemical analysis of *Myrtus communis* plant: Conventional versus microwave assisted-extraction procedures

N Bouaoudia-Madi, L Boulekbache-Makhlouf, N Kadri... - Journal of Complementary and Integrative Medicine, 2017

Keywords: antioxidant activity; microwave-assisted extraction; *Myrtus communis*; phenolic compounds