

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/230647659>

4-Carboxyanilinium hydrogensulfate

Article in *Acta Crystallographica Section E Structure Reports Online* · May 2007

DOI: 10.1107/S160053680701272X/dn2144sup2.hkl

CITATIONS

6

READS

61

5 authors, including:



Nourredine Benali-Cherif

University of Science and Technology Houari Boumediene

172 PUBLICATIONS 530 CITATIONS

[SEE PROFILE](#)



Amani Direm

Abbes Laghrour - Khenchela University

59 PUBLICATIONS 84 CITATIONS

[SEE PROFILE](#)



Lila Boukli

Abou Bakr Belkaid University of Tlemcen

34 PUBLICATIONS 78 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Intermolecular interactions in proton transfer compounds [View project](#)



Crystal engennering [View project](#)

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Editors: **W. Clegg** and **D. G. Watson**

4-Carboxyanilinium hydrogensulfate

Nourredine Benali-Cherif, Amani Direm, Fatima Allouche, Lila Boukli-H-Benmenni and Kawther Soudani

Copyright © International Union of Crystallography

Author(s) of this paper may load this reprint on their own web site provided that this cover page is retained. Republication of this article or its storage in electronic databases or the like is not permitted without prior permission in writing from the IUCr.

Nourredine Benali-Cherif,^{a*}
 Amani Direm,^a
 Fatima Allouche,^a
 Lila Boukli-H-Benmenni^b and
 Kawther Soudani^a

^aLaboratoire des Structures, Propriétés et Interactions Inter Atomiques (LASPI²A), Centre Universitaire de Khenchela, 40000 Khenchela, Algeria, and ^bDépartement de Chimie, Faculté des Sciences, Université Abou-Bekr Belkaid, Tlemcen, Algeria

Correspondence e-mail:
 benalicherif@hotmail.com

Key indicators

Single-crystal X-ray study
 $T = 293\text{ K}$
 Mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$
 R factor = 0.048
 wR factor = 0.123
 Data-to-parameter ratio = 19.4

For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.


4-Carboxyanilinium hydrogensulfate

The cohesion in the title structure (*p*-CPABS), $\text{C}_7\text{H}_8\text{NO}_2^+ \cdot \text{HSO}_4^-$, is assured by an intricate three-dimensional hydrogen-bonded network of types $\text{N}-\text{H} \cdots \text{O}$ and $\text{O}-\text{H} \cdots \text{O}$ involving anions, carboxyl and amino groups in addition to the ionic interactions.

Received 14 March 2007
 Accepted 18 March 2007

Comment

The electrical, magnetic and optical properties (Kagan *et al.*, 1999; Hill, 1998) of hybrid compounds (Mazeaud *et al.*, 2000; Soghomonian *et al.*, 1995; Mayer *et al.*, 1999; Bagieu-Beucher, 1990; Ravikumar *et al.*, 2002) make them very interesting materials because of their structural topologies and potential applications in the field of new materials science (Siegel *et al.*, 1998; Baker *et al.*, 1992).



p-Carboxyaniline (4-aminobenzoic acid, PABA) is an important biological molecule, acting as an antagonist to the action of the drug sulfonamide in competition for essential growth metabolites (Pauling & Hayward, 1964), as well as being an essential bacterial cofactor involved in the synthesis of folic acid (Robinson, 1966) and proving a particularly versatile reagent for structure extension through linear




Figure 1
 The asymmetric unit of *p*-CPABS, with the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level and H atoms are shown as small spheres of arbitrary radius. The hydrogen bond is shown as a double dashed line.




Figure 2
A partial packing view of (I), showing the hydrogen-bonding network. Hydrogen bonds are shown as dashed lines. H atoms not involved in hydrogen bonds have been omitted for clarity. [Symmetry codes: (i) $x, y - 1, z$; (ii) $1 - x, 1 - y, 1 - z$.]

hydrogen-bonding associations, through both the carboxylic acid and amine functional groups. This property was recognized as a possible tool for promoting co-crystallization, with the aim of designing noncentrosymmetric organic materials (Etter & Frankenbach, 1989).

In our systematic investigation of organic–inorganic hybrid materials, including sulfuric acid and various nucleic acids or amino acids, three structures have been already reported, namely diglycinium sulfate (Cherouana *et al.*, 2002), guanidinium sulfate monohydrate (Cherouana, Benali-Cherif *et al.*, 2003) and *m*-carboxyphenylammonium bisulfate (Cherouana, Bendjeddou *et al.*, 2003). We report here the fourth such compound, the title compound, (I).

The asymmetric unit of (I) (*p*-CPABS) contains one *p*-carboxyphenylammonium cation and one bisulfate anion (Fig. 1). The average of the terminal S–O bond lengths [1.4445 (14) Å] is shorter than the S–OH bond length [1.5609 (14) Å], which confirms the presence of the H atom in the bisulfate anion, supported also by its involvement in a hydrogen bond (see below). The O–S–O bond angles of the HSO₄[−] anions are in the range 102.63 (8)–115.35 (10)°, confirming a tetrahedral configuration, and are similar to those in other reported sulfates. The geometry of the organic cation is normal and in good agreement with that observed in *p*-carboxyphenylammonium dihydrogenmonophosphate monohydrate (Benali-Cherif *et al.*, 2002).

The three H atoms of the phenylammonium group are involved in extensive N–H···O hydrogen-bonding interactions with the O-atom acceptors of four different bisulfate anions (Table 1). In addition, there are secondary structure extensions involving the carboxylic acid group of the cations and the bisulfate anions (Table 1). Such an intricate three-dimensional hydrogen-bonding framework ensures the cohesion of the crystal structure (Fig. 2).

Experimental

Colourless prismatic single crystals of *p*-CPABS were obtained by slow evaporation, at room temperature, of an equimolar solution of 4-aminobenzoic acid (PABA) and sulfuric acid.

Crystal data

$C_7H_8NO_2^+ \cdot HO_4S^-$	$\gamma = 92.785 (6)^\circ$
$M_r = 235.21$	$V = 463.51 (10) \text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 5.2058 (6) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 7.5770 (7) \text{ \AA}$	$\mu = 0.36 \text{ mm}^{-1}$
$c = 11.837 (2) \text{ \AA}$	$T = 293 \text{ K}$
$\alpha = 93.356 (7)^\circ$	$0.25 \times 0.1 \times 0.05 \text{ mm}$
$\beta = 95.187 (7)^\circ$	

Data collection

Nonius KappaCCD area-detector diffractometer	2693 independent reflections
Absorption correction: none	2205 reflections with $I > 2\sigma(I)$
7756 measured reflections	$R_{\text{int}} = 0.091$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$	139 parameters
$wR(F^2) = 0.123$	H-atom parameters constrained
$S = 1.08$	$\Delta\rho_{\text{max}} = 0.41 \text{ e \AA}^{-3}$
2693 reflections	$\Delta\rho_{\text{min}} = -0.70 \text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
N1–H13···O6	0.89	2.13	2.918 (2)	147
N1–H12···O5 ⁱ	0.89	1.89	2.748 (2)	163
N1–H13···O4 ⁱⁱ	0.89	2.53	3.056 (2)	119
N1–H11···O4 ⁱⁱⁱ	0.89	1.92	2.802 (2)	170
O2–H2···O6 ^{iv}	0.82	1.85	2.647 (2)	164
O3–H3···O1 ^v	0.82	1.87	2.688 (2)	173

Symmetry codes: (i) $x, y - 1, z$; (ii) $-x + 1, -y + 1, -z + 2$; (iii) $-x + 2, -y + 1, -z + 2$; (iv) $-x + 2, -y + 1, -z + 1$; (v) $-x + 1, -y + 1, -z + 1$.

The OH and NH₃ H atoms of the anion and cations could be located in difference Fourier syntheses, but they were introduced in calculated positions and treated as riding on their parent atoms, with N–H = 0.89 Å and O–H = 0.82 Å, and with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{N}, \text{O})$. Aromatic H atoms were located in difference Fourier syntheses but were treated as riding on their parent C atoms, with C–H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Data collection: *KappaCCD Server Software* (Nonius, 1998); cell refinement: *DENZO* and *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* and *SCALEPACK*; program(s) used to solve structure: *SIR2004* (Burla *et al.*, 2005); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2003); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors thank Dr M. Giorgi, Faculté des Sciences et Techniques de Saint Jérôme, Marseille, France, for providing diffraction facilities, and the Centre Universitaire de Khenchela for financial support.

References

- Bagieu-Beucher, M. (1990). *Acta Cryst.* **C46**, 238–240.
- Baker, L.-J., Bowmaker, G. A., Healy, P. C., Skelton, B. W. & White, A. H. (1992). *J. Chem. Soc. Dalton Trans.* pp. 989–998.
- Benali-Cherif, N., Abouimrane, A., Sbai, K., Merazig, H., Cherouana, A. & Bendjeddou, L. (2002). *Acta Cryst.* **E58**, o160–o161.

- Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.
- Cherouana, A., Benali-Cherif, N. & Bendjeddou, L. (2003). *Acta Cryst.* **E59**, o180–o182.
- Cherouana, A., Benali-Cherif, N., Bendjeddou, L. & Merazig, H. (2002). *Acta Cryst.* **E58**, o1351–o1353.
- Cherouana, A., Bendjeddou, L. & Benali-Cherif, N. (2003). *Acta Cryst.* **E59**, o1790–o1792.
- Etter, M. C. & Frankenbach, G. A. (1989). *Chem. Mater.* **1**, 10–12.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Hill, C. L. (1998). *Chem. Rev.* **98**, 1–2.
- Kagan, C. R., Mitzi, D. B. & Dimitrakopoulos, C. D. (1999). *Science*, **286**, 945–947.
- Mayer, C. R., Herson, P. & Thouvenot, R. (1999). *Inorg. Chem.* **38**, 6152–6158.
- Mazeaud, A., Dromzee, Y. & Thouvenot, R. (2000). *Inorg. Chem.* **39**, 4735–4740.
- Nonius (1998). *KappaCCD Server Software*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Pauling, L. & Hayward, R. (1964). *The Architecture of Molecules*, p. 56. San Francisco: W. H. Freeman.
- Ravikumar, B. S., Sridhar, B. & Rajaram, R. K. (2002). *Acta Cryst.* **E58**, o879–o881.
- Robinson, F. A. (1966). *The Vitamin Co-factors of Enzyme Systems*, pp. 541–662. London: Pergamon.
- Sheldrick, G. M. (1997). *SHELXL97*. University of Göttingen, Germany.
- Siegel, R. K. O., Freisinger, E., Metzger, S. & Lippert, B. (1998). *J. Am. Chem. Soc.* **120**, 12000–12007.
- Soghomonian, V., Chen, Q., Haushalter, R. C. & Zubieta, J. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 223–226.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.