

Synthesis of Some New 3-Pyrrolidinylquinoline Derivatives via 1,3-Dipolar Cycloaddition of Stabilized Azomethine Ylides to Quinoliny α,β - Unsaturated Ketones

Abdelmalek Bouraiou, Abdelmadjid Debache, Salah Rhouati, Ali Belfaitah, N. Benali-Cherif, Bertrand Carboni

► **To cite this version:**

Abdelmalek Bouraiou, Abdelmadjid Debache, Salah Rhouati, Ali Belfaitah, N. Benali-Cherif, et al.. Synthesis of Some New 3-Pyrrolidinylquinoline Derivatives via 1,3-Dipolar Cycloaddition of Stabilized Azomethine Ylides to Quinoliny α,β - Unsaturated Ketones. Open Organic Chemistry Journal, Bentham Open, 2010, 4, pp.1-7. hal-00869747

HAL Id: hal-00869747

<https://hal.archives-ouvertes.fr/hal-00869747>

Submitted on 4 Oct 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Synthesis of Some New 3-Pyrrolidinylquinoline Derivatives via 1,3-Dipolar Cycloaddition of Stabilized Azomethine Ylides to Quinoliny α,β -Unsaturated Ketones

A. Bouraiou¹, A. Debache¹, S. Rhouati¹, A. Belfaitah*¹, N. Benali-Cherif² and B. Carboni³

¹Laboratoire des Produits Naturels d'Origine Végétale et de Synthèse Organique, Faculté des Sciences Exactes, Campus de Chaabat Ersas, Université Mentouri-Constantine, 25000, Algeria

²Laboratoire des Structures, Propriétés et Interactions Inter Atomiques (LASPI²A), Centre Universitaire de Khenchela, 40000 Khenchela, Algérie

³Sciences Chimiques de Rennes, UMR 6226 CNRS-Université de Rennes 1, Bat 10 A, Campus de Beaulieu, 35042 Rennes CEDEX, France

Abstract: N-Metallated azomethine ylide generated from methyl (*E*)-*N*-benzylideneglycinate, LiBr and triethylamine underwent cycloaddition to quinolyl α,β -unsaturated ketones with excellent diastereoselectivity to afford new functionalised 3-pyrrolidinylquinoline derivatives.

Keywords: 1,3-dipolar cycloaddition, quinoline, pyrrolidine synthesis, azomethine ylides.

INTRODUCTION

Quinolines derivatives have attracted considerable interest for many years due to their presence in the skeleton of a large number of bioactive compounds and natural products [1]. For example, quinoline alkaloids, such as quinine, chloroquine, mefloquine and amodiaquine, are used as efficient drugs for the treatment of malaria [2].

On the other hand 1,3-dipolar cycloaddition reactions of azomethine ylides with olefinic dipolarophiles had resulted in a number of novel heterocyclic scaffolds which are particularly useful for the creation of diverse chemical libraries of drug-like molecules for biological screening [3]. Functionalized pyrrolidine containing compounds are also of significant importance because of their biological activities and widespread employment in catalysis [4].

The coupling of this chemical entity with quinoline unit might as well be envisioned to bring with some biological activities. In this context, some limited investigations have been carried out which involved the combination of the quinolyl moiety and the pyrrolidine unit.

As a part of our program related to the preparation and biological evaluation of quinolyl derivatives [5], we have recently described a practical and an efficient synthesis of some 3-pyrrolidinylquinoline derivatives from quinoliny α,β -unsaturated esters as starting materials via 1,3-dipolar cycloaddition [6]. In a continuation of our efforts in this area, we report here an efficient procedure for the preparation of new pyrrolidine derivatives bearing a quinoline ring at C-3,

aroyl or acetyl group at C-4, and a phenyl substituent at C-5 via an 1,3-dipolar cycloaddition reaction of a stabilized metallo-azomethine ylide to quinoliny α,β -unsaturated ketones [7].

RESULTS AND DISCUSSION

Starting from the corresponding 2-chloro-3-formylquinoline derivatives **1**, chalcones **2a-2h** were synthesized by Claisen-Schmidt condensation reactions of appropriately substituted acetophenone in ethanol in the presence of 10% of aqueous NaOH [8]. The methylketone derivatives **2i-2j** were prepared from the aldehydes **1** via a Wittig reaction using methyl(triphenylphosphoranylidene) acetate and were obtained in good yields (Scheme 1).


The *E*-configured dipolarophiles (**2a-2j**) reacted with azomethine ylide, generated from methyl (*E*)-*N*-benzylideneglycinate in the presence of LiBr and triethylamine at room temperature, employing dry THF as the solvent (Scheme 2).

In accordance with literature reports [9, 10], this 1,3-dipolar cycloaddition reaction of the *in situ* generated metallo-azomethine ylide, exhibited high regio and stereoselectivity leading to the expected *syn-endo* cycloadduct (**3a-3j**).


All results reported below shown that pyrrolidines were obtained with conservation of the stereochemistry of starting alkenes [11], giving only one diastereoisomer with no evidence of any other isomers in the ¹H NMR spectra of the crude products (Table 1).

The structure of compounds **3a-3j** has been established by analogy and by comparison of their ¹H NMR with those reported [12]. The shielding of the protons of the methyl connected to the aroyl group attached at C-4 by the adjacent 5-phenyl ring ($\delta=1.83$ ppm) confirms the regiochemistry and demonstrated the 4,5-*cis* configuration relationship [6, 13].

*Address correspondence to this author at the Laboratoire des Produits Naturels d'Origine Végétale et de Synthèse Organique, Faculté des Sciences Exactes, Campus de Chaabat Ersas, Université Mentouri-Constantine, 25000, Algeria; Tel/Fax: 00 213 (0)31 81 88 62; E-mail: abelbelfaitah@yahoo.fr



Scheme 1. Synthesis of chalcone and methyl(vinylquinoline) ketone derivatives.



Scheme 2. Synthesis of quinolylypyrrolidine N-H derivatives (**3a-3j**).

Table 1. Synthesis of 3-pyrrolidinylquinolines **3**

Compound	R ¹	R ²	R ³	R ⁴	R	Yield (%)
3a	H	H	H	Me	2-MeC ₆ H ₄	70
3b	H	H	H	H	2-MeC ₆ H ₄	58
3c	OMe	H	H	OMe	2-MeC ₆ H ₄	71
3d	H	Me	H	H	2-MeC ₆ H ₄	62
3e	H	H	H	Me	4-OMeC ₆ H ₄	60
3f	H	OMe	H	H	4-OMeC ₆ H ₄	65
3g	H	H	H	Me	3,4-diMeOC ₆ H ₃	53
3h	H	H	H	Me	3,4,5-triMeOC ₆ H ₂	54
3i	H	Me	H	H	Me	67
3j	H	H	H	H	Me	59

The structure of compound **3a**, as representative example, was elucidated by detailed NMR studies (Table 2). The ¹H and ¹³C NMR assignments were made on the basis of high-field one and two-dimensional methods (HSQC, COSY, and NOESY H, H). The [2,4,5-*cis*] configuration of these pyrrolidines was confirmed by the observed NOE en-

hancement between the two pairs (H-2 and H-5) and (H-2 and H-4) (Table 2).

X-ray crystallography of **3e** (Fig. 1) showed an asymmetric unit which contains two independent molecules and the analysis demonstrate that the two stereoisomers have for each one, the absolute stereochemistry (2*S*,3*R*,4*S*,5*R*) and

Table 2. Significant ¹H, ¹³C NMR Data, Selected H-H Coupling NOE for **3a**

	δ ¹³ C	δ ¹ H (m, J)	¹ H{ ¹ H}n.O.e ^a	¹ H, ¹ H cosy
H-2	66.5	4.40 (d, 7.9)	H-5, H-4 Qui, 2-CO ₂ Me	4.69
H-5	66.4	4.96 (d, 8.1)	H-2, H-6 Ar, H-2 Ar	4.77
H-4	60.8	4.77 (t, 8.0)	H-2, H-4 Qui	4.69, 4.96
H-3	50.5	4.69 (t, 8.0)	H-4 Qui	4.40, 4.77

^aObtained by 2D-NOESY spectroscopy.

(2*R*,3*S*,4*R*,5*S*) of the new stereocenters created in the cycloaddition reactions [14].




Fig. (1). ORTEP of asymmetric unit of compound **3e** which contains two independent molecules projection down (010).

CONCLUSIONS

In conclusion, we report herein an efficient approach to 3-pyrrolidinylquinoline derivatives that exploits [3+2] cycloaddition reactions of azomethine ylides. This approach allows a diverse range of compounds to be generated in good yield and the pharmacological actions of the new pyrrolidine derivatives will be inspected afterwards.

EXPERIMENTAL SECTION

General Information

THF was freshly distilled from sodium/benzophenone, POCl₃ and CH₂Cl₂ from P₂O₅, DMF was kept for few hours over CaCl₂ and distilled from CaO and DME from NaH. EtOH was distilled from magnesium. Melting points were determined on an Electrothermal Digital Melting Points Apparatus IA 9200 and are uncorrected. IR spectra were performed on Shimadzu FT IR-8201 PC spectrophotometer and Perkin Elmer Spectrum One (FT-IR) spectrophotometer with a universal ATR sampling accessory. NMR spectra were recorded in CDCl₃ on a Brüker Avance DPX250 or Brüker Avance DMX300 spectrometer. Chemical shifts (δ) are given in ppm and J values in Hertz (Hz). column chromatography was performed on Merck silica gel (60, particle size 0.063-0.2 mm) using CHCl₃ or CH₂Cl₂ as eluent. Thin layer chromatography (TLC) was carried out on precoated Merck silica gel aluminium sheets 60 F₂₅₄. HRMS data were obtained on spectrometer MAT 311 (Centre Régional de Mesures Physiques de l'Ouest). X-Ray crystallographic analysis was performed with an Enraf-Nonius KAPPA CCD at 293 K using Mo K α radiation ($\lambda = 0.71073 \text{ \AA}$).

Substituted 2-chloroquinolyl-3-carbaldehydes have been synthesized according to reported methods [15]. Methyl benzylidene aminoacetate is obtained by treatment of benzaldehyde with methyl glycinate hydrochloride in basic medium [16].

General Method for the Synthesis of Chalcone Derivatives (2a-2h)

To a solution of 10% NaOH (520 mg, 13 mmol) in 95% ethanol (20 mL) was added 500 mg (2.61 mmol) of 2-chloro-3-formylquinoline and the acetophenone derivative (1.0 eq., 2.61 mmol). The mixture was stirred at 25 °C for 24 h. The contents were then cooled and poured into cold water then neutralized with dilute HCl. The solid obtained was filtered, washed, and dried on air to afford the crude chalcone.

(E)-3-(2-Chloro-8-methylquinolin-3-yl)-1-o-tolylprop-2-en-1-one (2a)

Yd 94%. *R_f* (CH₂Cl₂): 0.60. Mp 165-167 °C. IR ν_{max} (KBr) 1645 cm⁻¹ (C=O, ketone). ¹H NMR (250 MHz, CDCl₃) δ 8.48 (s, 1H), 8.00 (d, *J*=16.1, 1H), 7.75 (d, *J*=8.9, 1H), 7.52 (dd, *J*=9.1, *J*=2.4, 1H), 7.50-7.43 (m, 3H), 7.40 (ddd, *J*=8.9, *J*=8.1, *J*=2.3, 1H), 7.38 (dd, *J*=9.4, *J*=1.2, 1H), 7.25 (d, *J*=16.1, 1H), 2.75 (s, 3H), 2.50 (s, 3H). ¹³C NMR (62.5 MHz, CDCl₃) δ 195.2 (C=O), 149.1 (C), 147.1 (C), 140.5 (CH), 138.2 (C), 137.4 (C), 136.7 (CH), 136.4 (CH), 131.7 (C), 131.5 (CH), 130.9 (CH), 130.2 (C), 128.4 (CH), 127.4 (CH), 127.3 (C), 127.0 (CH), 125.8 (CH), 125.5 (CH), 20.4 (CH₃), 17.7 (CH₃). HRMS (EI): *m/z* [*M*⁺] Calcd. for C₂₀H₁₆NO³⁵Cl: 321.09204, found 321.0928.

(E)-3-(2-Chloroquinolin-3-yl)-1-o-tolylprop-2-en-1-one (2b)

Yd 88%. *R_f* (CH₂Cl₂): 0.65. Mp 120-124 °C. IR ν_{max} (KBr) 1685 cm⁻¹ (C=O, ketone). ¹H NMR (250 MHz, CDCl₃) δ 8.50 (s, 1H), 7.95 (d, *J*=15.9, 1H), 7.79 (dd, *J*=8.7, *J*=8.4, 1H), 7.52 (dd, *J*=8.9, *J*=2.4, 1H), 7.50-7.32 (4H, m), 7.37 (dd, *J*=8.4, *J*=1.2, 1H), 7.29 (d, *J*=15.9, 1H), 6.82 (td, *J*=9.1, *J*=2.1, 1H), 2.50 (s, 3H). ¹³C NMR (62.5 MHz, CDCl₃) δ 195.2 (C=O), 150.2 (C), 147.8 (C), 140.1 (CH), 138.1 (C), 137.5 (C), 137.3 (CH), 136.1 (CH), 133.4 (CH), 132.0 (CH), 131.6 (CH), 131.4 (CH), 131.3 (C), 130.6 (CH), 129.9 (CH), 127.2 (C), 125.5 (CH), 124.8 (CH), 20.4 (CH₃). HRMS (EI): *m/z* [*M*⁺] Calcd. for C₁₉H₁₄NO³⁵Cl: 307.07639, found 307.0769.

(E)-3-(2-Chloro-5,8-dimethoxyquinolin-3-yl)-1-o-tolylprop-2-en-1-one (2c)

Yd 87%. *R_f* (CH₂Cl₂): 0.62. Mp 169-170 °C. IR ν_{max} (KBr) 1672 cm⁻¹ (C=O, ketone). ¹H NMR (300 MHz, CDCl₃) δ 8.90 (s, 1H), 7.97 (d, *J*=16.0, 1H), 7.60 (d, *J*=7.5, 1H), 7.49 (d, *J*=7.4, 1H), 7.19-7.42 (m, 3H), 7.05 (d, *J*=16.0, 1H), 6.89 (d, *J*=8.6, 1H), 4.04 (s, 3H), 4.00 (s, 3H), 2.51 (s, 3H). ¹³C NMR (75.4 MHz, CDCl₃) δ 197.5 (C=O), 151.3 (C), 148.9 (C), 148.4 (C), 140.6 (CH), 138.7 (C), 135.6 (CH), 132.1 (C), 131.5 (C), 131.5 (CH), 130.9 (CH), 130.4 (C), 128.4 (CH), 127.9 (CH), 125.5 (CH), 121.8 (C), 109.8 (CH), 104.8 (CH), 56.2 (OCH₃), 55.8 (OCH₃), 20.4 (CH₃). HRMS (EI): *m/z* [*M*⁺] Calcd. for C₂₁H₁₈NO₃³⁵Cl: 367.09752, found 367.0961.

(E)-3-(2-Chloro-6-methylquinolin-3-yl)-1-o-tolylprop-2-en-1-one (2d)

Yield 91%. *R_f* (CH₂Cl₂): 0.59. Mp 143-145 °C. IR ν_{max} (KBr) 1654 cm⁻¹ (C=O, ketone). ¹H NMR (250 MHz, CDCl₃) δ 8.41 (s, 1H), 7.90 (d, *J*=16.0, 1H), 7.85 (d, *J*=8.3, 1H), 7.60-7.55 (m, 5H), 7.32 (d, *J*=16.0, 1H), 7.00 (dd, *J*=8.4, *J*=2.2, 1H), 2.43 (s, 3H), 2.66 (s, 3H). ¹³C NMR (62.9 MHz, CDCl₃) δ 197.8 (C=O), 150.1 (C), 146.6 (C), 140.5

(CH), 138.2 (C), 136.0 (C), 136.5 (CH), 135.4 (CH), 131.7 (C), 131.5 (CH), 130.9 (CH), 130.2 (C), 128.4 (CH), 128.1 (CH), 127.4 (C), 127.0 (CH), 125.6 (CH), 125.2 (CH), 20.4 (CH₃), 18.1 (CH₃). HRMS (EI): m/z [M⁺] Calcd. for C₂₀H₁₆NO³⁵Cl: 321.09204, found 321.0928.

(E)-3-(2-Chloro-8-methylquinolin-3-yl)-1-(4-methoxyphenyl)prop-2-en-1-one (2e)

Yield 97%. R_f (CH₂Cl₂): 0.55. Mp 131-134 °C. IR ν_{max}(KBr) 1662 cm⁻¹ (C=O, ketone). ¹H NMR (250 MHz, CDCl₃) δ 8.50 (s, 1H), 8.15 (d, J=15.7, 1H), 8.14 (d, J=8.8, 2H), 7.77-7.67 (m, 3H), 7.60 (d, J=15.7, 1H), 7.00 (d, J=8.8, 2H), 3.96 (s, 3H), 2.75 (s, 3H). ¹³C NMR (62.9 MHz, CDCl₃) δ 187.7 (C=O), 163.6 (C), 149.7 (C), 149.3 (C), 147.0 (C), 138.7 (CH), 136.6 (CH), 136.3 (C), 131.5 (CH), 131.0 (CH), 130.5 (C), 130.3 (CH), 127.7 (2×CH), 127.3 (CH), 127.0 (C), 113.9 (2×CH), 55.5 (OCH₃), 17.7 (CH₃). HRMS (EI): m/z [M⁺] Calcd. for C₂₀H₁₆NO₂³⁵Cl: 337.08696, found 337.0869.

(E)-3-(2-Chloro-6-methoxyquinolin-3-yl)-1-(4-methoxyphenyl)prop-2-en-1-one (2f)

Yield 90%. R_f (CH₂Cl₂): 0.65. Mp 142-145 °C. IR ν_{max}(KBr) 1653 cm⁻¹ (C=O, ketone). ¹H NMR (250 MHz, CDCl₃) δ 8.56 (s, 1H), 8.21 (d, J=15.9, 1H), 7.57 (d, J=8.1, 1H), 7.55 (d, J=8.7, 2H), 7.22 (d, J=8.9, 1H), 7.15 (d, J=15.9, 1H), 7.12 (s, 1H), 6.75 (d, J=8.6, 2H), 4.00 (s, 3H), 3.73 (s, 3H). ¹³C NMR (75.4 MHz, CDCl₃) δ 195.8 (C=O), 162.5 (C), 155.4 (C), 142.7 (C), 142.5 (C), 137.9 (C), 134.4 (CH), 132.6 (C), 131.5 (CH), 131.8 (CH), 127.0 (CH), 126.2 (C), 125.9 (2×CH), 121.6 (CH), 113.1 (2×CH), 108.6 (CH), 55.5 (OCH₃), 54.4 (OCH₃). HRMS (EI): m/z [M⁺] Calcd. for C₂₀H₁₆NO₃³⁵Cl: 353.08187, found 353.0825.

(E)-3-(2-Chloro-8-methylquinolin-3-yl)-1-(3,4-dimethoxyphenyl)prop-2-en-1-one (2g)

Yield 90%. R_f (CH₂Cl₂): 0.64. Mp 183-184 °C. IR ν_{max}(KBr) 1658 cm⁻¹ (C=O, ketone). ¹H NMR (250 MHz, CDCl₃) δ 8.50 (s, 1H), 8.25 (d, J=15.7, 1H), 7.75-7.67 (m, 4H), 7.52 (d, J=15.7, 1H), 7.50 (s, 1H), 7.00 (d, J=8.4, 1H), 4.00 (s, 3H), 3.96 (s, 3H), 2.78 (s, 3H). ¹³C NMR (62.9 MHz, CDCl₃): δ 195.5 (C=O), 153.5 (C), 150.0 (C), 149.2 (C), 146.9 (C), 145.7 (CH), 138.7 (CH), 136.6 (C), 136.4 (CH), 136.3 (CH), 131.5 (C), 130.8 (CH), 128.6 (C), 127.7 (CH), 127.3 (C), 124.9 (CH), 110.7 (CH), 109.8 (CH), 56.2 (OCH₃), 56.1 (OCH₃), 17.7 (CH₃). HRMS (EI): m/z [M⁺] Calcd. for C₂₁H₁₈NO₃³⁵Cl: 367.09752, found 367.0997.

(E)-3-(2-Chloro-8-methylquinolin-3-yl)-1-(3,4,5-trimethoxyphenyl)prop-2-en-1-one (2h)

Yield 72 %. R_f (CH₂Cl₂): 0.65. Mp 122-124 °C. IR ν_{max}(KBr) 1656 cm⁻¹ (C=O, ketone). ¹H NMR (250 MHz, CDCl₃) δ 8.49 (s, 1H), 8.20 (d, J=15.5, 1H), 7.89 (d, J=7.1, 1H), 7.40 (d, J=7.4, 1H), 7.34 (d, J=7.1, 1H), 7.09 (d, J=15.5, 1H), 6.89 (s, 2H), 3.97 (s, 3H), 3.93 (s, 3H), 3.83 (s, 6H), 2.65 (s, 3H); ¹³C NMR (62.9 MHz, CDCl₃) δ 195.0 (C=O), 151.7 (C), 148.9 (C), 144.3 (C), 143.5 (C), 136.3 (C), 136.2 (CH), 136.0 (C), 132.4 (C), 131.3 (CH), 130.4 (C), 128.9 (CH), 128.1 (CH), 127.4 (CH), 127.3 (C), 125.4 (CH), 105.6 (2×CH), 59.9 (OCH₃), 55.9 (OCH₃), 54.9 (OCH₃), 17.8 (CH₃). HRMS (EI): m/z [M⁺] Calcd. for C₂₂H₂₀NO₄³⁵Cl: 367.09752, found 367.0997.

General Procedure for the Preparation of Methyl(vinylquinoline) Ketone Derivatives

A suspension of the ylide Ph₃P=CHCOMe (313 mg, 1.1 mmol) and the 2-chloro-3-formylquinoline (191.5 mg, 1.0 mmol) in DME (10 mL) was refluxing for three hours. After cooling to room temperature, the mixture was filtered and the filtrate was condensed under reduced pressure. The residue was then purified by column chromatography over silica gel (CH₂Cl₂) to give the olefinic product.

(E)-4-(2-Chloro-6-methylquinolin-3-yl)but-3-en-2-one (2i)

Yield 56%. R_f (CH₂Cl₂): 0.75. Mp 73-74 °C. IR ν_{max}(KBr) 1645 cm⁻¹ (C=O, ketone). ¹H NMR (250 MHz, CDCl₃) δ 8.25 (s, 1H), 7.90 (d, J=16.3, 1H), 7.73 (s, 1H), 7.57-7.54 (m, 2H), 6.75 (d, J=16.3, 1H), 2.52 (s, 3H), 2.45 (s, 3H); ¹³C NMR (62.9 MHz, CDCl₃) δ 197.8 (C=O), 149.1 (C), 146.5 (C), 138.1 (CH), 137.9 (C), 135.4 (CH), 134.0 (CH), 130.8 (CH), 128.0 (CH), 127.1 (C), 127.0 (C), 126.8 (CH), 27.3 (CH₃), 21.6 (CH₃). HRMS (EI): m/z [M⁺] Calcd. for C₁₄H₁₂NO³⁵Cl: 245.06074, found 245.0607.

(E)-4-(2-Chloroquinolin-3-yl)but-3-en-2-one (2j)

Yield 70%. R_f (CH₂Cl₂): 0.72. Mp 175-177 °C. IR ν_{max}(KBr) 1651 cm⁻¹ (C=O, ketone). ¹H NMR (300 MHz, CDCl₃) δ 8.42 (s, 1H), 8.04 (d, 8.4, 1H), 7.98(d, J=16.3, 1H), 7.88 (d, J=8.2, 1H), 7.80 (t, J=7.9, 1H), 7.62 (t, J=7.3, 1H), 6.82 (d, J=16.3, 1H), 2.48 (s, 3H); ¹³C NMR (75.4 MHz, CDCl₃) δ 197.6 (C), 148.5 (C), 138.0 (CH), 136.1 (CH), 133.4 (C), 131.9 (C), 131.7 (CH), 131.0 (C), 128.4 (CH), 128.0 (CH), 127.7 (C), 126.4 (CH), 30.9 (CH₃). HRMS (EI): m/z [M⁺] Calcd. for C₁₃H₁₀NO³⁵Cl: 231.04509, found 231.0452.

General Procedure for the Preparation of Quinolylypyrrolidine N-H Derivatives

To 1.5 eq. of lithium bromide dissolved in dry THF (e.g. 0.5 g in 40 mL) was added, under magnetic stirring and at room temperature, 1 eq. of benzylidene glycine imine, 1 eq. of substituted quinolyl α,β-unsaturated ketone derivative and 1.2 eq. of dry Et₃N. The reaction mixture was kept under stirring, at room temperature and the progress of the reaction was monitored by TLC still disappearance of starting product. The mixture was diluted with ether (15 mL) and work up by treatment with saturated aqueous ammonium chloride (10 mL). The organic layers were separated and dried over anhydrous MgSO₄. The filtrate was concentrated under reduced pressure and the residue was subjected to column chromatography on silica gel using CHCl₃ as eluent to afford pure product.

Methyl 4-(2-methylbenzoyl)-3-(2-chloro-8-methylquinolin-3-yl)-5-phenylpyrrolidine-2-carboxylate (3a)

Yield 70%. R_f (CHCl₃): 0.34. Mp 74-76 °C. IR (ATR) ν 3332, 2953, 1737, 1678, 1574, 1479, 1455, 1372, 1335, 1239, 753 cm⁻¹. ¹H NMR (300 MHz, CDCl₃) δ 8.26 (s, 1H), 7.69 (d, J=8.1, 1H), 7.57 (d, J=6.9, 1H), 7.46 (m, 2H), 7.27 (td, J=7.5, J=1.3, 1H), 7.16-7.02 (m, 6H), 6.99 (d, J=7.5, 1H), 4.96 (d, J=8.1, 1H), 4.77 (t, J=8.0, 1H), 4.69 (t, J=8.0, 1H), 4.40 (d, J=7.9, 1H), 3.80 (s, 3H), 3.00 (brs, 1H), 2.80 (s, 3H), 1.83 (s, 3H). ¹³C NMR (75.4 MHz, CDCl₃) δ 202.0 (C=O), 173.3 (C=O), 150.0 (C), 146.3 (C), 139.8 (C), 139.0 (C), 137.9 (C), 137.8 (CH), 136.9 (C), 132.4 (C), 132.2

(CH), 131.8 (CH), 130.9 (CH), 129.0 (CH), 128.7 (2xCH), 128.1 (CH), 127.8 (C), 127.7 (2xCH), 127.5 (CH), 125.7 (CH), 125.6 (CH), 66.5 (CH, C-2), 66.4 (CH, C-5), 60.8 (CH, C-4), 52.9 (OCH₃), 50.5 (CH, C-3), 21.0 (CH₃), 18.1 (CH₃). MS (ESI): m/z 499.1 (MH⁺, 100), 439 (10), 394 (3), 322 (30), 178 (77), 119 (21), 91 (5), 60 (5).

Methyl 4-(2-methylbenzoyl)-3-(2-chloroquinolin-3-yl)-5-phenylpyrrolidine-2-carboxylate (3b)

Yield 58%. R_f (CHCl₃): 0.34. Mp 79-81 °C. IR (ATR) v 3347, 2926, 1735, 1676, 1567, 1488, 1454, 1331, 1202, 1133, 731 cm⁻¹. ¹H NMR (300 MHz, CDCl₃) δ 8.31 (s, 1H), 8.04 (d, J=8.5, 1H), 8.86 (d, J=8.1, 1H), 7.72 (td, J=7.7, J=1.1, 1H), 7.59 (t, J=7.4, 1H), 7.46 (d, J=7.7, 1H), 7.24 (t, J=7.6, 1H), 7.17-7.11 (m, 6H), 6.99 (d, J=7.5, 1H), 4.96 (d, J=8.2, 1H), 4.81 (t, J=8.1, 1H), 4.70 (t, J=8.1, 1H), 4.37 (d, J=8.3, 1H), 3.79 (s, 3H), 2.88 (brs, 1H), 1.83 (s, 3H). ¹³C NMR (75.4 MHz, CDCl₃) δ 201.7 (C=O), 173.2 (C=O), 151.3 (C), 147.1 (C), 139.9 (C), 139.0 (C), 138.4 (C), 137.8 (C), 137.5 (CH), 132.8 (C), 132.2 (CH), 131.9 (CH), 130.9 (CH), 129.0 (CH), 128.7 (2xCH), 128.6 (CH), 128.6 (CH), 128.1 (CH), 127.7 (2xCH), 127.7 (CH), 125.8 (CH), 66.6 (CH, C-2), 66.3 (CH, C-5), 61.0 (CH, C-4), 53.0 (OCH₃), 50.2 (CH, C-3), 21.0 (CH₃). MS (ESI): m/z 485.2 (MH⁺, 100), 425 (8), 380 (5), 308 (28), 178 (75), 146 (7), 119 (56), 91 (8), 60(7). HRMS (ESI): m/z [MH⁺] Calcd. for C₂₉H₂₆N₂O₃³⁵Cl: 485.16320, found 485.1628.

Methyl 4-(2-methylbenzoyl)-3-(2-chloro-5,8-dimethoxyquinolin-3-yl)-5-phenylpyrrolidine-2-carboxylate (3c)

Yield 71%. R_f (CHCl₃): 0.34. Mp 153-154 °C. IR (ATR) v 3339, 2953, 1733, 1674, 1617, 1592, 1481, 1330, 1263, 1014, 725 cm⁻¹. ¹H NMR (300 MHz, CDCl₃) δ 8.66 (s, 1H), 7.39 (d, J=7.7, 1H), 7.24 (td, J= 7.4, J=1.2, 1H), 7.15-7.11 (m, 6H), 6.98 (m, 2H), 6.78 (d, J=8.5, 1H), 4.96 (d, J=8.1, 1H), 4.73 (t, J=7.9, 1H), 4.72 (t, J=8.0, 1H), 4.35 (d, J=8.2, 1H), 4.03 (s, 3H), 3.98 (s, 3H), 3.77 (s, 3H), 3.23 (brs, 1H), 1.80 (s, 3H). ¹³C NMR (75.4 MHz, CDCl₃) δ 202.14 (C=O), 173.2 (C=O), 151.3 (C), 148.9 (C), 148.6 (C), 139.7 (C), 139.1 (C), 139.0 (C), 138.0 (C), 132.7 (C), 132.1 (CH), 131.8 (CH), 128.9 (CH), 128.7 (2xCH), 128.0 (CH), 127.7 (2xCH), 127.4 (CH), 125.7 (CH), 121.1 (C), 108.6 (CH), 104.9 (CH), 67.1 (CH, C-2), 66.5 (CH, C-5), 61.0 (CH, C-4), 56.5 (OCH₃), 56.1 (OCH₃), 52.9 (OCH₃), 51.0 (CH, C-3), 20.9 (CH₃). MS (ESI): m/z 545.1 (MH⁺, 100), 509 (10), 485 (5), 368 (38), 332 (3), 178 (38), 119 (15), 91 (1), 60 (1). HRMS (ESI): m/z [MH⁺] Calcd. for C₃₁H₃₀N₂O₅³⁵Cl: 545.18432, found 545.1839.

Methyl 4-(2-methylbenzoyl)-3-(2-chloro-6-methylquinolin-3-yl)-5-phenylpyrrolidine-2-carboxylate (3d)

Yield 62%. R_f (CHCl₃): 0.29. Mp 83-85 °C. IR (ATR) v 3336, 2952, 1736, 1677, 1597, 1494, 1338, 1215, 1129, 1044, 731 cm⁻¹. ¹H NMR (300 MHz, CDCl₃) δ 8.21 (s, 1H), 7.93 (d, J=8.6, 1H), 7.62 (s, 1H), 7.57 (dd, J=8.7, J=1.7, 1H), 7.43 (d, J=7.7, 1H), 7.26 (td, J= 7.4, J=1.2, 1H), 7.17-7.14 (m, 6H), 6.99 (d, J=7.5, 1H), 4.95 (d, J=8.0, 1H) 4.78 (t, J=8.0, 1H), 4.67 (t, J=8.1, 1H), 4.37 (d, J=7.9, 1H), 3.79 (s, 3H), 2.78 (brs, 1H), 2.55 (s, 3H), 1.83 (s, 3H); ¹³C NMR (75.4 MHz, CDCl₃) δ 201.8 (C=O), 173.3 (C=O), 150.3 (C), 145.7 (C), 139.8 (C), 139.0 (C), 137.8 (C), 137.8 (C), 136.9 (CH), 133.1 (CH), 132.6 (C), 132.2 (CH), 131.9 (CH), 129.0

(CH), 128.7 (2xCH), 128.3 (CH), 128.1 (CH), 127.8 (C), 127.7 (2xCH), 126.6 (CH), 125.7 (CH), 66.4 (CH, C-2), 66.3 (CH, C-5), 61.0 (CH, C-4), 52.9 (OCH₃), 50.2 (CH, C-3), 22.0 (CH₃), 21.0 (CH₃). HRMS (ESI): m/z [MH⁺] Calcd. for C₃₀H₂₈N₂O₃³⁵Cl: 499.17885, found 499.1793.

Methyl 4-(4-methoxybenzoyl)-3-(2-chloro-8-methylquinolin-3-yl)-5-phenylpyrrolidine-2-carboxylate (3e)

Yield 60%. R_f (CHCl₃): 0.32. Mp 108 °C. IR (ATR) v 3342, 2924, 1740, 1656, 1595, 1512, 1433, 1374, 1227, 1024, 749, 697 cm⁻¹. ¹H NMR (300 MHz, CDCl₃) δ 8.26 (s, 1H), 7.69 (d, J=8.1, 1H), 7.57-7.50 (m, 3H), 7.48 (t, J=7.7, 1H), 7.18-7.08 (m, 5H), 6.72 (d, J=8.8, 2H), 4.98 (d, J=8.1, 1H), 4.72 (t, J=7.9, 1H), 4.56 (t, J=7.9, 1H), 4.43 (d, J=8.0, 1H), 3.83 (s, 3H), 3.78 (s, 3H), 3.22 (brs, 1H), 2.78 (s, 3H); ¹³C NMR (75.4 MHz, CDCl₃) δ 197.6 (C=O), 173.6 (C=O), 163.6 (C), 150.3 (C), 146.3 (C), 138.6 (C), 137.4 (CH), 136.8 (C), 132.9 (C), 130.8 (2xCH), 130.6 (C), 129.5 (CH), 128.6 (2xCH), 128.1 (CH), 127.7 (C), 127.6 (2xCH), 127.5 (CH), 125.7 (CH), 113.7 (2xCH), 66.8 (CH, C-2), 66.0 (CH, C-5), 58.8 (CH, C-4), 55.7 (OCH₃), 53.0 (OCH₃), 50.4 (CH, C-3), 18.14 (CH₃). MS (ESI): m/z 515.3 (MH⁺, 85), 455 (10), 410 (8), 338 (24), 178 (100), 135 (21), 118 (10), 91 (3), 60 (3). Calcd for C₃₀H₂₇N₂O₄Cl: C, 69.97; H, 5.28; N, 5.44. Found C, 69.33; H, 5.36; N, 5.39.

Methyl 4-(4-methoxybenzoyl)-3-(2-chloro-6-methoxyquinolin-3-yl)-5-phenylpyrrolidine-2-carboxylate (3f)

Yield 65%. R_f (CHCl₃): 0.28. Mp 100-103 °C. IR (ATR) v 3354, 2952, 1734, 1655, 1596, 1494, 1353, 1223, 1170, 907, 700 cm⁻¹. ¹H NMR (300 MHz, CDCl₃) δ 8.19 (s, 1H), 7.91 (d, J=9.0, 1H), 7.55 (d, J=8.8, 2H), 7.36 (dd, J=9.1, J=2.6, 1H), 7.18-7.05 (m, 6H), 6.70 (d, J=8.8, 2H), 4.97 (d, J=8.1, 1H), 4.68 (t, J=8.0, 1H), 4.40 (t, J=8.0, 1H), 4.39 (d, J=7.9, 1H), 3.93 (s, 3H), 3.83 (s, 3H), 3.78 (s, 3H), 3.10 (brs, 1H). ¹³C NMR (75.4 MHz, CDCl₃) δ 197.6 (C=O), 175.6 (C=O), 163.6 (C), 158.7 (C), 148.6 (C), 143.1 (C), 138.5 (C), 136.0 (CH), 133.4 (C), 130.8 (2xCH), 130.6 (C), 129.9 (CH), 128.8 (C), 128.6 (2xCH), 128.2 (CH), 127.6 (2xCH), 123.5 (CH), 113.7 (2xCH), 105.3 (CH), 66.7 (CH, C-2), 66.0 (CH, C-5), 58.6 (CH, C-4), 56.0 (OCH₃), 55.7 (OCH₃), 53.0 (OCH₃), 50.3 (CH, C-3). MS (ESI): m/z 531.3 (MH⁺, 73), 471 (7), 426 (7), 366 (5), 354 (28), 178 (100), 135 (24), 118 (17), 91 (8), 60 (7). HRMS (ESI): m/z [MH⁺] Calcd. for C₃₀H₂₈N₂O₅³⁵Cl: 531.16868, found 531.1680.

Methyl 4-(3,4-dimethoxybenzoyl)-3-(2-chloro-8-methylquinolin-3-yl)-5-phenylpyrrolidine-2-carboxylate (3g)

Yield 53%. R_f (CHCl₃): 0.30. Mp 115-116 °C. IR (ATR) v 3350, 1758, 1653, 1586, 1492, 1850, 709 cm⁻¹. ¹H NMR (300 MHz, CDCl₃) δ 8.29 (s, 1H), 7.64-7.00 (m, 10H), 6.71 (d, J=8.0, 1H), 5.00 (d, J=8.0, 1H), 4.78 (t, J=8.0, 1H), 4.60 (t, J=8.2, 1H), 4.53 (d, J=8.0, 1H), 3.87 (s, 3H), 3.85 (s, 3H), 3.76 (s, 3H), 2.78 (s, 3H). ¹³C NMR (75.4 MHz, CDCl₃) δ 197.2 (C=O), 173.2 (C=O), 153.0 (C), 150.0 (C), 148.6 (C), 145.8 (C), 137.9 (CH), 36.9 (C), 136.4 (C), 133.4 (C), 130.4 (CH), 129.6 (C), 128.2 (2xCH), 127.8 (CH), 127.3 (C), 127.2 (2xCH), 127.1 (CH), 125.3 (CH), 122.9 (CH), 110.0 (CH), 109.5 (CH), 66.2 (CH), 65.4 (CH), 58.4 (CH), 55.9 (OCH₃), 55.7 (OCH₃), 52.6 (OCH₃), 49.1 (CH), 17.7 (CH₃). HRMS (ESI): m/z [MH⁺] Calcd. for C₃₁H₃₀N₂O₅³⁵Cl: 545.18433, found 545.1843.

Methyl 4-(3,4,5-trimethoxybenzoyl)-3-(2-chloro-8-methylquinolin-3-yl)-5-phenyl pyrrolid-ine-2-carboxylate (3h)

Yield 54%. R_f (CHCl₃): 0.38. Mp 209-210 °C. IR (ATR) ν 3354, 2949, 1732, 1648, 1494, 1250, 1185, 915, 708 cm⁻¹. ¹H NMR (300 MHz, CDCl₃) δ 8.28 (1H, s), 7.71 (d, $J=7.9$, 1H), 7.59 (d, $J=7.9$, 1H), 7.48 (t, $J=7.9$, 1H), 7.16-7.10 (m, 5H), 6.77 (s, 2H), 4.95 (d, $J=8.1$, 1H), 4.71 (t, $J=8.0$, 1H), 4.42-4.44 (m, 2H), 3.87 (s, 3H), 3.83 (s, 3H), 3.76 (s, 6H), 3.30 (s, 3H), 2.78 (s, 3H). ¹³C NMR (75.4 MHz, CDCl₃) δ 198.0 (C=O), 173.4 (C=O), 152.5 (C), 149.8 (C), 145.9 (C), 142.4 (C), 137.9 (C), 136.8 (CH), 136.4 (C), 132.4 (C), 132.2 (C), 130.5 (CH), 128.3 (2 \times CH), 128.0 (CH), 127.3 (2 \times CH), 127.2 (C), 127.2 (CH), 125.3 (CH), 105.8 (2 \times CH), 66.4 (CH), 65.3 (CH), 60.8 (OCH₃), 58.8 (CH), 56.2 (OCH₃), 52.7 (OCH₃), 50.0 (CH), 17.7 (CH₃). HRMS (ESI): m/z [MH⁺] Calcd. for C₃₂H₃₂N₂O₆³⁵Cl : 575.19489, found 575.1917.

Methyl 4-(methylketone)-3-(2-chloro-6-methylquinolin-3-yl)-5-phenylpyrrolidine-2-carboxylate (3i)

Yellow oil. Yield 67%. R_f (CHCl₃): 0.54. IR (ATR) ν 3350, 2957, 1732, 1657, 1285, 1226, 1085, 910, 695 cm⁻¹. ¹H NMR (300 MHz, CDCl₃) δ 8.24 (s, 1H), 7.94 (d, $J=8.5$, 1H), 7.68 (s, 1H), 7.60 (dd, $J=8.5$, $J=1.6$, 1H), 7.37-7.25 (m, 5H), 4.87 (d, $J=8.0$, 1H), 4.53 (t, $J=8.1$, 1H), 4.35 (d, $J=7.9$, 1H), 3.82 (s, 3H), 3.60 (dd, $J=7.9$, $J=4.8$, 1H), 2.55 (s, 3H), 1.50 (s, 3H). ¹³C NMR (75.4 MHz, CDCl₃) δ 208.0 (C=O), 172.6 (C=O), 149.6 (C), 145.0 (C), 137.4 (C), 136.8 (CH), 135.9 (C), 132.8 (CH), 132.2 (C), 128.8 (2 \times CH), 128.4 (CH), 127.5 (CH), 127.3 (C), 126.9 (2 \times CH), 126.4 (CH), 64.8 (CH), 64.3 (CH), 63.9 (CH), 52.7 (OCH₃), 49.0 (CH), 28.3 (OCH₃), 21.4 (CH₃). HRMS (EI): m/z [M⁺] Calcd. for C₂₄H₂₂N₂O₃³⁵Cl: 421.13190, found 421.1329.

Methyl 4-(methylketone)-3-(2-chloroquinolin-3-yl)-5-phenylpyrrolidine-2-carboxylate (3j)

Yellow oil. Yield 59%. R_f (CHCl₃): 0.45. IR (ATR) ν 3367, 1739, 1651, 1342, 1237, 1110 cm⁻¹. ¹H NMR (300 MHz, CDCl₃) δ 8.29 (s, 1H), 8.08 (d, $J=8.4$, 1H), 7.90 (d, $J=8.1$, 1H), 7.73 (t, $J=7.0$, 1H), 7.61 (t, $J=7.1$, 1H), 7.30-7.45 (m, 5H), 4.85 (d, $J=8.0$, 1H), 4.51 (t, $J=8.0$, 1H), 4.32 (d, $J=7.9$, 1H), 3.83 (s, 3H), 3.61 (dd, $J=8.0$, $J=4.8$, 1H). ¹³C NMR (75.4 MHz, CDCl₃) δ 207.8 (C=O), 172.8 (C=O), 150.6 (C), 146.6 (C), 139.2 (C), 137.3 (CH), 136.7 (CH), 132.7 (C), 130.6 (CH), 128.9 (2 \times CH), 128.4 (CH), 128.1 (CH), 127.5 (CH), 127.3 (C), 127.0 (2 \times CH), 65.2 (CH), 65.1 (CH), 52.7 (OCH₃), 49.2 (CH), 31.2 (CH), 28.3 (CH₃). HRMS (EI): m/z [M⁺] Calcd. for C₂₃H₂₁N₂O₃³⁵Cl: 408.12407, found 408.1221.

ACKNOWLEDGEMENTS

ANDRS (Agence Nationale pour le Développement de la Recherche en Santé) and MESRES (Ministère de l'Enseignement Supérieur et de la Recherche Scientifique) are gratefully acknowledged for their financial support. AB thanks l'Agence Universitaire de la Francophonie (AUF) for financial support.

REFERENCES

- [1] (a) Padwa, A.; Brodney, M. A.; Liu, B.; Satake, K.; Wu, T. *J. Org. Chem.* **1999**, *64*, 3595. (b) Egan, T. J. *J. Inorg. Biochem.* **2006**, *100*, 916. (c) Sahu, N. P.; Pal, C.; Mandal, N. B.; Banerjee, S.; Raha, M.; Kundu, A. P.; Basu, A.; Ghosh, M.; Roy, K.; Bandyopadhyay, S. *Bioorg. Med. Chem.* **2002**, *10*, 1687.
- [2] (a) Robert, A.; Meunier, B. *Chem. Soc. Rev.* **1998**, *27*, 273. (b) Dow, G. S.; Heady, T. N.; Bhattacharjee, A. K.; Caridha, D.; Gerena, L.; Gettayacamin, M.; Lanteri, C. A.; Obaldia, N.; Roncal, N.; Shearer, T.; Smith, P. L.; Tungtaeng, A.; Wolf, L.; Cabezas, M.; Yourick, D.; Smith, K. S. *Antimicrob. Chemother.* **2006**, *50*, 4132. (c) Tjitra, E.; Baker, J.; Suprianto, S.; Cheng, Q.; Anstey, N. M. *Antimicrob. Agents Chemother.* **2002**, *46*, 3947. (d) Sowunmi, A.; Sowunmi, C. O.; Adedeji, A. A.; Oduold, A. M. *Clin. Drug Invest.* **2001**, *21*, 33.
- [3] (a) Tsuge, O.; Kanemasa, S. *Advances in Heterocyclic Chemistry*, Katritzky, A. R., Ed.; Academic: San Diego, **1989**; Vol. *45*, p. 231. (b) Gohelf, K. V. In *Cycloaddition Reactions in Organic Synthesis* Kobayashi, S.; Jørgensen, K. A.; Eds. Wiley-VCH, Weinheim, **2002**, Chapter 6, pp. 211-247.
- [4] For some recent references on this subject see: (a) Fraley, M. E.; Hartman, G. D.; Hungate, R. W. WO 0162252, August 30, 2001. (b) Arrington, K. L.; Bilodeau, M. T.; Fraley, M.; Hartman, G. D.; Hoffman, W. F.; Hungate, R.; Kim, Y. WO 0129025, April 26, 2001. (c) Fraley, M. E.; Hambaugh, S. R.; Hungate, R. W. WO 0128993, April 26, 2001. (d) Witherup, K.; Ranson, R. W.; Graham, A. C.; Barnard, A. M.; Salvatore, M. J.; Limma, W. C.; Anderson, P. S.; Pitzengerger, S. M.; Varga, S. L. *J. Am. Chem. Soc.* **1995**, *117*, 6682. (e) Kravchenko, D. V.; Kysil, V. M.; Tkachenko, S. E.; Maliarchouk, S.; Okun Ilya, M. I.; Ivachtchenko, A. V. *Eur. J. Med. Chem.* **2005**, *40*, 1377. (f) Alcaide, B and Almdendros, P. *Angew. Chem. Int. Ed.* **2003**, *42*, 858.
- [5] (a) Bouraiou, A.; Menasra, H.; Debache, A.; Rhouati, S.; Belfaitah, A. *J. Soc. Alger. Chim.* **2006**, *16*, 171. (b) Kedjadja, A.; Moussaoui, F.; Debache, A.; Rhouati, S.; Belfaitah, A. *J. Soc. Alger. Chim.* **2004**, *14*, 225.
- [6] Bouraiou, A.; Debache, A.; Rhouati, S.; Carboni, B.; Belfaitah, A. *J. Heterocycl. Chem.* **2008**, *45*, 329.
- [7] For a similar approach using azomethine ylides generated from N-alkylamino acids and formaldehyde, see: Menasra, H.; Kedjadja, A.; Rhouati, S.; Carboni, B.; Belfaitah, A. *Synth. Commun.* **2005**, *35*, 2779.
- [8] (a) Rezig, R.; Chebah, M.; Rhouati, S.; Ducki, S.; Lawrence, N. *J. Soc. Alger. Chim.* **2000**, *10*, 111. (b) Moussaoui, F.; Belfaitah, A.; Debache, A.; Rhouati, S. *J. Soc. Alger. Chim.* **2002**, *12*, 71.
- [9] (a) Tsuge, O.; Kanemasa, S.; Yoshioka, M. *J. Org. Chem.* **1988**, *53*, 1384. (b) Butler, R. N.; Farrell, D. M. *J. Chem. Res. (S)*. **1998**, *2*, 82. (c) Annunziata, R.; Cinquina, M.; Cozzi, M.; Raimondi, L.; Pilati, T. *Tetrahedron Asymmetry* **1991**, *2*, 1329, and references cited therein.
- [10] (a) Lown, J. W. In *3-Dipolar Cycloaddition Chemistry* Padwa, A., Ed. Wiley, New York, **1984**, Vol. *1*, p. 653. (b) Coldham, L.; Collis, A. J.; Mould, R. J.; Robinson, D. E. *Synthesis* **1995**, *9*, 1147. (c) Wittland, C.; Arend, M.; Rish, N. *Synthesis* **1996**, *3*, 367. (d) Galley, G.; Liebscher, J.; Pätz, M. *J. Org. Chem.* **1995**, *60*, 5005. (e) Nyerges, M.; Rudas, M.; Tóth, G.; Herényi, B.; Bitter, I.; Töke, L. *Tetrahedron* **1995**, *51*, 13321.
- [11] (a) Grigg, R.; Montgomery, J.; Somasunderam, A. *Tetrahedron* **1992**, *48*, 10431. (b) Pak, C. S.; Nyerges, M. *Bull. Korean Chem. Soc.* **1999**, *20*, 633.
- [12] (a) Pak, C. S.; Nyerges, M. *Synlett* **1999**, *8*, 1271. (b) Nyerges, M.; Rudas, M.; Bitter, I.; Töke, L. *Tetrahedron* **1997**, *53*, 3269. (c) Fejes, I.; Töke, L.; Blaskó, G.; Nyerges, M.; Pak, C. S. *Tetrahedron* **2000**, *56*, 8545.
- [13] Belfaitah, A.; Isly, M.; Carboni, B. *Tetrahedron Lett.* **2004**, *45*, 1969.
- [14] The single-crystal growth was carried out in CHCl₃ at room temperature. X-ray diffraction data have been collected with an Enraf Nonius KAPPA CCD at 293 K using Mo K α radiation ($\lambda = 0.71073$ Å). The crystal **3e** belong to the triclinic space group P_1 , with unit-cell parameters $a = 12.4982(2)$ Å, $b = 13.6473(4)$ Å, $c = 18.3062(2)$ Å, $\alpha = 110.709(3)^\circ$, $\beta = 105.647(4)^\circ$ and $\gamma = 89.9690(10)^\circ$. The asymmetric unit contains two molecules. Bond angles, bond distances and other crystallographic data (excluding structure factors) for the structures reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary Publication No. CCDC-662754 for compound **3e**. Copies of the data can be obtained free of charge on application to the CCDC, 12 Union Road, Cambridge, CB2 1EZ, UK.

- [15] (a) Meth-Cohn, O.; Rhouati, S.; Tarnowski, B.; Robinson, A. J. *Chem. Soc. Perkin Trans.* **1981**, *1*, 1537. (b) Tóth, J.; Blaskó, G.; Dancsó, A.; Tóke, L.; Nyerges, M. *Synth. Commun.* **2006**, 3581.
- [16] Stork, G.; Leong, A. W.; Touzin, A. M. *J. Org. Chem.* **1976**, *41*, 3491.

Received: December 14, 2009

Revised: February 27, 2010

Accepted: March 12, 2010

© Bouraiou *et al.*; Licensee Bentham Open.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.